

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 March 2001 (22.03.2001)

PCT

(10) International Publication Number
WO 01/19845 A1

- (51) International Patent Classification⁷: C07H 21/04, C12Q 1/68
- (21) International Application Number: PCT/US00/25479
- (22) International Filing Date:
14 September 2000 (14.09.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/398,522 15 September 1999 (15.09.1999) US
- (71) Applicant: THE JOHNS HOPKINS UNIVERSITY
SCHOOL OF MEDICINE [US/US]; Suite 906, 111
Market Street, Baltimore, MA 21202 (US).
- (72) Inventor: ISSA, Jean-Pierre; 112310 Longworth Lane,
Houston, TX 77024 (US).
- (74) Agent: HAILE, Lisa, A.; Gray Cary Ware & Freidenrich
LLP, Suite 1600, 4365 Executive Drive, San Diego, CA
92121-2189 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/19845 A1

(54) Title: CACNA1G POLYNUCLEOTIDE, POLYPEPTIDE AND METHODS OF USE THEREFOR

(57) Abstract: A novel T-type calcium channel (CACNA1G) is provided, as are polynucleotides encoding the same. CACNA1G has been implicated in cellular proliferative disorders. More specifically, it has been observed that the methylation state of specific regions within CpG islands associated with the CACNA1G gene correlates with a number of cancerous phenotypes involving a variety of tissue and cell types. Also provided are methods for detecting cellular proliferative disorders by determining the methylation state of genes or regulatory regions associated therewith, including CACNA1G, as well as kits containing reagents for performing invention methods.

CACNA1G POLYNUCLEOTIDE , POLYPEPTIDE and METHODS OF USE THEREFOR**STATEMENT AS TO FEDERALLY SPONSORED RESEARCH**

5

This invention was made with Government support under Grant No. CA43318 and CA54396, awarded by the National Cancer Institute and Grant No. CA43318, a Colon Cancer Spore Grant. The government may have certain rights in the invention.

10

FIELD OF THE INVENTION

The present invention relates generally to the regulation of gene expression and more specifically to a method of determining the DNA methylation status of CpG sites in a given locus and correlating the methylation status with the presence of a cell proliferative disorder.

15

BACKGROUND OF THE INVENTION

DNA methylases transfer methyl groups from the universal methyl donor S-adenosyl methionine to specific sites on the DNA. Several biological functions have been attributed to the methylated bases in DNA. The most established biological function for methylated DNA is the protection of DNA from digestion by cognate restriction enzymes. The restriction modification phenomenon has, so far, been observed only in bacteria. Mammalian cells, however, possess a different methylase that exclusively methylates cytosine residues that are 5' neighbors of guanine (CpG). This modification of cytosine residues has important regulatory effects on gene expression, especially when involving CpG rich areas, known as CpG islands, located in the promoter regions of many genes.

20

25

30

Methylation has been shown by several lines of evidence to play a role in gene activity, cell differentiation, tumorigenesis, X-chromosome inactivation, genomic imprinting and other major biological processes (Razin, A., H., and Riggs, R.D. eds.

in DNA Methylation Biochemistry and Biological Significance, Springer-Verlag, New York, 1984). In eukaryotic cells, methylation of cytosine residues that are immediately 5' to a guanosine, occurs predominantly in CG poor regions (Bird, A., *Nature*, 321:209, 1986). In contrast, CpG islands remain unmethylated in normal
5 cells, except during X-chromosome inactivation (Migeon, et al., *supra*) and parental specific imprinting (Li, et al., *Nature*, 366:362, 1993) where methylation of 5' regulatory regions can lead to transcriptional repression. De novo methylation of the Rb gene has been demonstrated in a small fraction of retinoblastomas (Sakai, et al., *Am. J. Hum. Genet.*, 48:880, 1991), and recently, a more detailed analysis of the VHL
10 gene showed aberrant methylation in a subset of sporadic renal cell carcinomas (Herman, et al., *Proc. Natl. Acad. Sci., U.S.A.*, 91:9700, 1994). Expression of a tumor suppressor gene can also be abolished by de novo DNA methylation of a normally unmethylated CpG island (Issa, et al., *Nature Genet.*, 7:536, 1994; Herman, et al., *supra*; Merlo, et al., *Nature Med.*, 1:686, 1995; Herman, et al., *Cancer Res.*, 56:722,
15 1996; Graff, et al., *Cancer Res.*, 55:5195, 1995; Herman, et al., *Cancer Res.*, 55:4525, 1995).

Human cancer cells typically contain somatically altered nucleic acid, characterized by mutation, amplification, or deletion of critical genes. In addition,
20 the nucleic acid from human cancer cells often displays somatic changes in DNA methylation (E.R. Fearon, et al., *Cell*, 61:759, 1990; P.A. Jones, et al., *Cancer Res.*, 46:461, 1986; R. Holliday, *Science*, 238:163, 1987; A. De Bustros, et al., *Proc. Natl. Acad. Sci., USA*, 85:5693, 1988); P.A. Jones, et al., *Adv. Cancer Res.*, 54:1, 1990; S.B. Baylin, et al., *Cancer Cells*, 3:383, 1991; M. Makos, et al., *Proc. Natl. Acad. Sci., USA*, 89:1929, 1992; N. Ohtani-Fujita, et al., *Onco-gene*, 8:1063, 1993). However,
25 the precise role of abnormal DNA methylation in human tumorigenesis has not been established.

Aberrant methylation of normally unmethylated CpG islands has been described as a frequent event in immortalized and transformed cells, and has been
30 associated with transcriptional inactivation of defined tumor suppressor genes in

human cancers. In the development of colorectal cancers (CRC), a series of tumor suppressor genes (TSG) such as APC, p53, DCC and DPC4 are inactivated by mutations and chromosomal deletions (reviewed in Kinzler and Vogelstein 1996). Some of these alterations result from a chromosomal instability phenotype described in a subset of CRC (Lengauer et al., 1997a). Recently, an additional pathway has been shown to be involved in a familial form of CRC, hereditary non-polyposis colorectal cancer. The cancers from these patients show a characteristic mutator phenotype which causes microsatellite instability (MI), and mutations at other gene loci such as TGF-beta-RII (Markowitz et al., 1995) and BAX (Rampino et al., 1997). This phenotype usually results from mutations in the mismatch repair (MMR) genes hMSH2 and hMLH1 (reviewed by Peltomaki, and de la Chapelle, 1997). A subset of sporadic CRC also show MI, but mutations in MMR genes appear to be less frequent in these tumors (Liu et al., 1995; Moslein et al., 1996).

Another molecular defect described in CRC is CpG island (CGI) methylation. CGIs are short sequences rich in the CpG dinucleotide and can be found in the 5' region of about half of all human genes. Methylation of cytosine within 5' CGIs is associated with loss of gene expression and has been seen in physiological conditions such as X chromosome inactivation and genomic imprinting (reviewed in Latham, 1996). Aberrant methylation of CGIs has been detected in genetic diseases such as the fragile-X syndrome, in aging cells and in neoplasia. About half of the tumor suppressor genes which have been shown to be mutated in the germline of patients with familial cancer syndromes have also been shown to be aberrantly methylated in some proportion of sporadic cancers, including Rb, VHL, p16, hMLH1, and BRCA1 (reviewed in Baylin, S.B., Herman, J.G., Graff, J.R., Vertino, P.M. and Issa, J.P. Alterations in DNA methylation: a fundamental aspect of neoplasia, *Adv. Cancer Res.* 72:141-196 1998). TSG methylation in cancer is usually associated with (1) lack of gene transcription and (2) absence of coding region mutation. Thus it has been proposed that CGI methylation serves as an alternative mechanism of gene inactivation in cancer.

The causes and global patterns of CGI methylation in human cancers remain poorly defined. Aging could play a factor in this process because methylation of several CGIs could be detected in an age-related manner in normal colon mucosa as well as in CRC (Issa, J.P., Vertino, P.M., Boehm, C.D., Newsham, I.F. and Baylin, S.B. Methylation of the oestrogen receptor CpG island links ageing and neoplasia in human colon. *Nat Genet.* 7:536-540, 1994). In addition, aberrant methylation of CGIs has been associated with the MI phenotype in CRC as well as specific carcinogen exposures (Issa et al., 1996) supra. However, an understanding of aberrant methylation in CRC has been somewhat limited by the small number of CGIs analyzed to date. In fact, previous studies have suggested that large numbers of CGIs are methylated in immortalized cell lines (Antequera, F., Boyes, J. and Bird, A. High levels of de novo methylation and altered chromatin structure at CpG islands in cell lines. *Cell* 62:503-514, 1990), and it is not well understood whether this global aberrant methylation is caused by the cell culture conditions or whether they are an integral part of the pathogenesis of cancer.

Most of the methods developed to date for detection of methylated cytosine depend upon cleavage of the phosphodiester bond alongside cytosine residues, using either methylation-sensitive restriction enzymes or reactive chemicals such as hydrazine which differentiate between cytosine and its 5-methyl derivative. Genomic sequencing protocols which identify a 5-MeC residue in genomic DNA as a site that is not cleaved by any of the Maxam Gilbert sequencing reactions have also been used, but still suffer disadvantages such as the requirement for large amount of genomic DNA and the difficulty in detecting a gap in a sequencing ladder which may contain bands of varying intensity.

Mapping of methylated regions in DNA has relied primarily on Southern hybridization approaches, based on the inability of methylation-sensitive restriction enzymes to cleave sequences which contain one or more methylated CpG sites. This method provides an assessment of the overall methylation status of CpG islands,

including some quantitative analysis, but is relatively insensitive and requires large amounts of high molecular weight DNA.

Another method utilizes bisulfite treatment of DNA to convert all
5 unmethylated cytosines to uracil. The altered DNA is amplified and sequenced to show the methylation status of all CpG sites. However, this method is technically difficult, labor intensive and without cloning amplified products, it is less sensitive than Southern analysis, requiring approximately 10% of the alleles to be methylated for detection.

10

Identification of the earliest genetic changes in tumorigenesis is a major focus in molecular cancer research. Diagnostic approaches based on identification of these changes are likely to allow implementation of early detection strategies and novel therapeutic approaches targeting these early changes might lead to more effective
15 cancer treatment.

SUMMARY OF THE INVENTION

The present invention is based on the finding that several genes are newly
20 identified as being differentially methylated in cancer. This seminal discovery is useful for cancer screening, risk-assessment, prognosis, minimal-residual disease identification, staging and identification of therapeutic targets. The identification of new genes that are methylated in cancer, aging or diseases associated with aging increases the likelihood of finding genes methylated in a particular cancer; increases
25 the sensitivity and specificity of methylation detection; allows methylation profiling using multiple genes; and allows identification of new targets for therapeutic intervention. The invention also provides a newly identified gene that is a target for hypermethylation in human tumors. This new gene, as well as genes newly identified as hypermethylated in cancer and aging or aging diseases provides markers which can

be used diagnostically, prognostically and therapeutically over the course of such disorders.

In a first embodiment, the invention provides a nucleic acid molecule
5 comprising a coding region for a T-type calcium channel, CACNA1G, and regulatory
sequences associated therewith. The discovery of CpG islands, and in particular,
methylated CpG islands in the region approximately 300-800 base pairs upstream
from the CACNA1G translation initiation start site, led to a method of the present
invention for correlating methylated CpG islands with various cancers. In one aspect
10 of this embodiment, the nucleic acid molecule encoding CACNA1G and the
associated regulatory sequences and CpG-rich regions include the nucleic acid
sequence set forth in SEQ ID NO:51 (Figure 3A). Also provided is a polypeptide
having an amino acid sequence as set forth in SEQ ID NO:52 and Figure 3B. The
methylation state of CpG islands in CACNA1G, associated regulatory regions, and
15 other genes is indicative of the presence of a cellular proliferative disorder in a subject
from which the CpG-containing nucleic acid is isolated.

In another embodiment, there are provided methods for detecting a cellular
proliferative disorder in a subject. The subject may have or be at risk of having a
20 cellular proliferative disorder. The method of the invention is useful for diagnostic as
well as prognostic analyses. One method for detecting a cellular proliferative disorder
in a subject includes contacting a nucleic acid-containing specimen from the subject
with an agent that provides a determination of the methylation state of at least one
gene or associated regulatory region of the gene; and identifying aberrant methylation
25 of regions of the gene or regulatory region, wherein aberrant methylation is identified
as being different when compared to the same regions of the gene or associated
regulatory region in a subject not having the cellular proliferative, thereby detecting a
cellular proliferative disorder in the subject. The method includes multiplexing by
utilizing a combination of primers for more than one loci, thereby providing a
30 methylation "profile" for more than one gene or regulatory region.

For the first time, the invention provides methylated forms of the following genes and/or their associated regulatory sequences: APOB, CACNA1G, CDX2, EGFR, FBN1, GPR37, HSPA6, IQGAP2, KL, PAR2, PITX2, PTCH, SDC1, and SDC4 (see Table 5). In addition, the invention provides the CpG-rich regions from these genes that are hypermethylated (see Figures 4A-4F (SEQ ID NO:105-119).

Invention methods include determining, in a nucleic acid-containing specimen taken from a subject, the methylation state of a gene or regulatory sequences associated therewith, wherein the expression or non-expression of the gene is associated with the presence of the cellular proliferative disorder, and identifying as having a cellular proliferative disorder a subject that has aberrant methylation of regions of the gene or associated regulatory sequences when compared to the same regions of the gene in a subject not having the cellular proliferative disorder. In one aspect of this embodiment, the methylated regions of the gene and associated regulatory sequences are contained within CpG islands (i.e., CpG rich regions). In particular, the aberrant methylation typically includes hypermethylation as compared with the same regions of the gene or regulatory sequences in a subject not having the cellular proliferative disorder.

Determining the methylation state of the gene includes contacting the nucleic acid-containing specimen with an agent that modifies unmethylated cytosine, amplifying a CpG-containing nucleic acid in the specimen by means of CpG-specific oligonucleotide primers, wherein the oligonucleotide primers distinguish between modified methylated and nonmethylated nucleic acid, and detecting the methylated nucleic acid based on the presence or absence of amplification products produced in said amplifying step. The method includes optionally contacting the amplification products with a methylation sensitive restriction endonuclease. Other methods for determining methylation status of a gene and/or regulatory sequences are well known in the art and are described more fully herein.

In another embodiment, the present invention provides a method of treating a cell proliferative disorder associated with CACNA1G or other methylated genes described herein, comprising administering to a subject with the disorder, a therapeutically effective amount of reagent which modulates CACNA1G or other methylated genes' expression. For example, since CACNA1G -associated disorders typically involve hypermethylation of CACNA1G polynucleotide sequence, a polynucleotide sequence which contains a non-methylatable nucleotide analog is utilized for treatment of a subject. Further, the invention provides a method of gene therapy comprising introducing into cells of a host subject, an expression vector comprising a nucleotide sequence encoding CACNA1G or other methylated genes described herein, in operable linkage with a promoter.

In another embodiment of the present invention there is provided a kit useful for the detection of a cellular proliferative disorder in a subject having or at risk for having a cellular proliferative disorder. Invention kits include a carrier means compartmentalized to receive a sample, one or more containers comprising a first container containing a reagent which modifies unmethylated cytosine and a second container containing primers for amplification of a CpG-containing nucleic acid, wherein the primers distinguish between modified methylated and nonmethylated nucleic acid, and optionally, a third container containing a methylation sensitive restriction endonuclease.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1A shows the relative positions of MINT31 and CACNA1G and associated CpG regions.

5 Figure 1B provides a magnified depiction of MINT31, CACNA1G and CpG rich regions 1-8.

Figure 2 depicts the correlation of methylation and the expression of CACNA1G.

Figures 3A and 3B show the nucleic acid sequence and deduced amino acid
10 sequence of CACNA1G (SEQ ID NO:51 and 52, respectively).

Figures 4A-4F show the CpG-rich regions of the genes depicted in Table 5 (SEQ ID NO:105-119).

Figure 5 is the nucleotide sequence of MINT31. (SEQ ID NO:120).

DETAILED DESCRIPTION OF THE INVENTION

15 It has been determined that an aberrant methylation state of nucleic acids in certain genes, particularly regulatory sequences, is diagnostic for the presence or potential development of a cellular proliferative disorder in subjects bearing the
20 aberrantly methylated nucleic acids. More particularly, the hypermethylation of certain nucleotides localized in CpG islands has been shown to affect the expression of genes associated with the CpG islands; typically such hypermethylated genes have reduced or abolished expression, primarily due to down-regulated transcription. Using a recently developed PCR-based technique called methylated CpG island
25 amplification (MCA), several nucleic acid molecules aberrantly methylated in a colon cancer cell line were identified. One DNA fragment, termed MINT31, mapped to human chromosome 17q21 where frequent loss of heterozygosity (LOH) has been detected in various human tumors. By characterizing the genomic sequence around this area, a gene encoding a T-type calcium channel, CACNA1G, was identified as a
30 target for hypermethylation in human tumors. Using RT-PCR, expression of CACNA1G was detected in normal colon and bone marrow, but expression was

absent in 5 tumor cell lines where methylation was found. After treatment with the methylation inhibitor 5-deoxy-azacytidine, the expression of CACNA1G was restored in all 5 cell lines. Detailed methylation mapping of the 5'CpG island by bisulfite-PCR revealed that methylation of a region 300 to 800 base pairs upstream of the translation initiation site closely correlated with the inactivation of CACNA1G. Aberrant methylation of CACNA1G was also examined in various human primary tumors, and was detected in 17 of 49 (35%) colorectal cancers, 4 of 16 (25%) gastric cancers, and 3 of 23 (13%) acute myelogenous leukemia cases. While not wanting to be bound by a particular theory, it is believed that inactivation of CACNA1G may play a role in cancer development by modulating calcium signaling, which potentially affects cell proliferation and apoptosis.

Thus, in one embodiment of the present invention, there are provided nucleic acids comprising the coding region for a T-type calcium channel and regulatory sequences associated therewith. Specifically, the T-type calcium channel and associated regulatory sequences comprise CACNA1G. In a more preferred embodiment, the CACNA1G is the human form of the gene. An exemplary CACNA1G gene and associated regulatory sequences is set forth in SEQ ID NO:51.

The invention provides methylated and unmethylated nucleic acid encoding CACNA1G (SEQ ID NO:51). Polynucleotides include DNA, cDNA and RNA sequences which encode CACNA1G polypeptide (SEQ ID NO:52). It is understood that naturally occurring, synthetic, and intentionally manipulated polynucleotides are included. For example, CACNA1G nucleic acid may be subject to site-directed mutagenesis, or the like. The nucleic acid sequences for CACNA1G also include antisense sequences, and sequences encoding dominant negative forms of CACNA1G, as well as sequences encoding functional fragments thereof. It is understood that naturally occurring, synthetic, and intentionally manipulated polynucleotides are included.

Methylated nucleic acid sequences are also provided. For the first time, the present invention provides methylated chemical structures for the following genes: APOB, CACNA1G, CDX2, EGFR, FBN1, GPR37, HSPA6, IQGAP2, KL, PAR2, PITX2, PTCH, SDC1, and SDC4. One of skill in the art can now readily locate the
5 CpG-rich sequences associated with these genes and identify such methylated forms of the genes/regulatory sequences by methods described herein (The gene sequences can be identified in a gene database found at <http://www.ncbi.nlm.nih.gov/UniGene/index.html>). The invention provides CpG-rich regions from the above genes as set forth in SEQ ID Nos:105-119. Thus, in yet
10 another embodiment, the invention provides an isolated nucleic acid molecule having at least one methylated Cytosine of a CpG dinucleotide in a CpG-rich region and encoding a gene selected from APOB, CACNA1G, CDX2, EGFR, FBN1, GPR37, HSPA6, IQGAP2, KL, PAR2, PITX2, PTCH, SDC1, and SDC4. The methylated C residue of a CpG dinucleotide is located within a CpG-rich region selected from SEQ
15 ID NO:105-118 and SEQ ID NO:119.

The polynucleotides of the invention include "degenerate variants" which are sequences that encode the same polypeptide yet vary in sequence as a result of the genetic code. There are 20 natural amino acids, most of which are specified by more
20 than one codon. Therefore, all degenerate nucleotide sequences are included in the invention as long as the amino acid sequence of a polypeptide encoded by the nucleotide sequence of SEQ ID NO:51 is functionally unchanged.

The term "polynucleotide" or "nucleic acid sequence" refers to a polymeric
25 form of nucleotides at least 10 bases in length. An "isolated polynucleotide" is a polynucleotide that is not immediately contiguous with both of the coding sequences with which it is immediately contiguous (one on the 5' end and one on the 3' end) in the naturally occurring genome of the organism from which it is derived. Thus, an isolated polynucleotide may include a coding region with its associated regulatory
30 sequences. The term therefore includes, for example, a recombinant DNA which is

incorporated into a vector; into an autonomously replicating plasmid or virus; or into the genomic DNA of a prokaryote or eukaryote, or which exists as a separate molecule (e.g., a cDNA) independent of other sequences. The nucleotides of the invention can be ribonucleotides, deoxyribonucleotides, or modified forms of either
5 nucleotide. Specifically, methylated forms of nucleotides in a polynucleotide sequence, such as regions 1-8 of CACNA1G as described herein, are also included. The term includes single and double forms of DNA.

As will be understood by those of skill in the art, when the sequence is RNA,
10 the deoxynucleotides A, G, C, and T of SEQ ID NO:51, are replaced by ribonucleotides A, G, C, and U, respectively. Also included in the invention are fragments of the above-described nucleic acid sequences that are at least 15 bases in length, which is sufficient to permit the fragment to selectively hybridize to DNA that encodes SEQ ID NO:52. The term "selectively hybridize" refers to hybridization
15 under moderately or highly stringent conditions (See, Maniatis, as cited herein) which excludes non-related nucleotide sequences.

The CACNA1G nucleic acid sequence includes the disclosed sequence and sequences that encode conservative variations of the polypeptides encoded by
20 CACNA1G polynucleotide provided herein. The term "conservative variation" as used herein denotes the replacement of an amino acid residue by another, biologically similar residue. Examples of conservative variations include the substitution of one hydrophobic residue such as isoleucine, valine, leucine or methionine for another, or the substitution of one polar residue for another, such as the substitution of arginine
25 for lysine, glutamic for aspartic acid, or glutamine for asparagine, and the like. The term "conservative variation" also includes the use of a substituted amino acid in place of an unsubstituted parent amino acid provided that antibodies raised to the substituted polypeptide also immunoreact with the unsubstituted polypeptide.

CACNA1G nucleic acid sequences can be expressed in vitro by DNA transfer into a suitable host cell. "Host cells" are cells in which a vector can be propagated and its DNA expressed. The cell may be prokaryotic or eukaryotic. The term also includes any progeny of the subject host cell. It is understood that all progeny may not be identical to the parental cell since there may be mutations that occur during replication. However, such progeny are included when the term "host cells" is used. Methods of stable transfer, meaning that the foreign DNA is continuously maintained in the host, are known in the art.

10 In one aspect, the CACNA1G nucleic acid sequences may be inserted into an expression vector. The term "expression vector" refers to a plasmid, virus or other vehicle known in the art that has been manipulated by insertion or incorporation of the sequence of interest genetic sequences. Polynucleotide sequence which encode sequence of interest can be operatively linked to expression control sequences.

15 "Operatively linked" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. An expression control sequence operatively linked to a coding sequence is ligated such that expression of the coding sequence is achieved under conditions compatible with the regulatory or expression control sequences. As used herein, the terms "regulatory sequences" and "expression control sequences" refers to nucleic acid sequences that regulate the expression of a nucleic acid sequence to which it is operatively linked.

20 Expression control sequences are operatively linked to a nucleic acid sequence when the expression control sequences control and regulate the transcription and, as appropriate, translation of the nucleic acid sequence. Thus expression control sequences can include appropriate promoters, enhancers, transcription terminators, a start codon (i.e., ATG) in front of a protein-encoding gene, splicing signal for introns, maintenance of the correct reading frame of that gene to permit proper translation of mRNA, and stop codons. The terms "regulatory sequences" and "expression control sequences" are intended to include, at a minimum, components whose presence can influence expression, and can also include additional components whose presence is

25

30

advantageous, for example, leader sequences and fusion partner sequences. An example of an expression control sequence includes a promoter.

A "promoter" is a minimal sequence sufficient to direct transcription. Also
5 included in the invention are those promoter elements which are sufficient to render promoter-dependent gene expression controllable for cell-type specific, tissue-specific, or inducible by external signals or agents; such elements may be located in the 5' or 3' regions of the gene. Both constitutive and inducible promoters, are included in the invention (see, e.g., Bitter et al., *Methods in Enzymology* 153:516-
10 544, 1987). For example, when cloning in bacterial systems, inducible promoters such as pL of bacteriophage plac, ptrp, ptac (ptrp-lac hybrid promoter) and the like may be used. When cloning in mammalian cell systems, promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the retrovirus long terminal repeat; the adenovirus late promoter; the
15 vaccinia virus 7.5K promoter) may be used. Promoters produced by recombinant DNA or synthetic techniques may also be used to provide for transcription of the nucleic acid sequences of the invention.

In the present invention, the CACNA1G polynucleotide sequence may be
20 inserted into an expression vector which contains a promoter sequence which facilitates the efficient transcription of the inserted genetic sequence of the host. The expression vector typically contains an origin of replication, a promoter, as well as specific genes which allow phenotypic selection of the transformed cells. Vectors suitable for use in the present invention include, but are not limited to the T7-based
25 expression vector for expression in bacteria (Rosenberg et al., *Gene* 56:125, 1987), the pMSXND expression vector for expression in mammalian cells (Lee and Nathans, *J. Biol. Chem.* 263:3521, 1988) and baculovirus-derived vectors for expression in insect cells. The DNA segment can be present in the vector operably linked to regulatory elements, for example, a promoter (e.g., T7, metallothionein I, or polyhedron
30 promoters).

CACNA1G polynucleotide sequences can be expressed in either prokaryotes or eukaryotes. Hosts can include microbial, yeast, insect and mammalian organisms. Methods of expressing DNA sequences having eukaryotic or viral sequences in prokaryotes are well known in the art. Biologically functional viral and plasmid DNA
5 vectors capable of expression and replication in a host are known in the art. Such vectors are used to incorporate DNA sequences of the invention.

"Transformation" means a genetic change induced in a cell following incorporation of new DNA (i.e., DNA exogenous to the cell). Where the cell is a
10 mammalian cell, the genetic change is generally achieved by introduction of the DNA into the genome of the cell (i.e., stable).

Thus, a "transformed cell" is a cell into which (or into an ancestor of which) has been introduced, by means of recombinant DNA techniques, a DNA molecule
15 encoding sequence of interest. Transformation of a host cell with recombinant DNA may be carried out by conventional techniques as are well known to those skilled in the art. Where the host is prokaryotic, such as *E. coli*, competent cells which are capable of DNA uptake can be prepared from cells harvested after exponential growth phase and subsequently treated by the CaCl_2 method using procedures well known in
20 the art. Alternatively, MgCl_2 or RbCl can be used. Transformation can also be performed after forming a protoplast of the host cell if desired.

When the host is a eukaryote, such methods of transfection of DNA as calcium phosphate co-precipitates, conventional mechanical procedures such as
25 microinjection, electroporation, insertion of a plasmid encased in liposomes, or virus vectors may be used. Eukaryotic cells can also be cotransformed with DNA sequences encoding the sequence of interest, and a second foreign DNA molecule encoding a selectable phenotype, such as the herpes simplex thymidine kinase gene. Another method is to use a eukaryotic viral vector, such as simian virus 40 (SV40) or bovine
30 papilloma virus, to transiently infect or transform eukaryotic cells and express the

protein (see for example, Eukaryotic Viral Vectors, Cold Spring Harbor Laboratory, Gluzman ed., 1982).

Methods which are well known to those skilled in the art can be used to
5 construct expression vectors containing the CACNA1G coding sequence and
appropriate transcriptional/translational control signals. These methods include in
vitro recombinant DNA techniques, synthetic techniques, and in vivo
recombination/genetic techniques. See, for example, the techniques described in
Maniatis, et al., 1989 Molecular Cloning A Laboratory Manual, Cold Spring Harbor
10 Laboratory, N.Y.

A variety of host-expression vector systems may be utilized to express the
CACNA1G coding sequence. These include but are not limited to microorganisms
such as bacteria transformed with recombinant bacteriophage DNA, plasmid DNA or
15 cosmid DNA expression vectors containing the CACNA1G coding sequence; yeast
transformed with recombinant yeast expression vectors containing the CACNA1G
coding sequence; plant cell systems infected with recombinant virus expression
vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or
transformed with recombinant plasmid expression vectors (e.g., Ti plasmid)
20 containing the CACNA1G coding sequence; insect cell systems infected with
recombinant virus expression vectors (e.g., baculovirus) containing the CACNA1G
coding sequence; or animal cell systems infected with recombinant virus expression
vectors (e.g., retroviruses, adenovirus, vaccinia virus) containing the CACNA1G
coding sequence, or transformed animal cell systems engineered for stable expression.
25 Since CACNA1G has not been confirmed to contain carbohydrates, both bacterial
expression systems as well as those that provide for translational and post-
translational modifications may be used; e.g., mammalian, insect, yeast or plant
expression systems.

30 Depending on the host/vector system utilized, any of a number of suitable
transcription and translation elements, including constitutive and inducible promoters,

transcription enhancer elements, transcription terminators, etc. may be used in the expression vector (see e.g., Bitter, et al., *Methods in Enzymology* 153:516-544, 1987). For example, when cloning in bacterial systems, inducible promoters such as pL of bacteriophage .gamma., plac, ptrp, ptac (ptrp-lac hybrid promoter) and the like may be
5 used. When cloning in mammalian cell systems, promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses(e.g., the retrovirus long terminal repeat; the adenovirus late promoter; the vaccinia virus 7.5K promoter) may be used. Promoters produced by recombinant DNA or synthetic techniques may also be used to provide for transcription of the inserted CACNA1G
10 coding sequence. In addition, the endogenous CACNA1G promoter may also be used to provide transcription machinery of CACNA1G .

In bacterial systems a number of expression vectors may be advantageously selected depending upon the use intended for the expressed. For example, when large
15 quantities of CACNA1G are to be produced, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Those which are engineered to contain a cleavage site to aid in recovering are preferred. Such vectors include but are not limited to the E. coli expression vector pUR278 (Ruther, et al., *EMBO J.* 2:1791, 1983), in which the CACNA1G coding sequence
20 may be ligated into the vector in frame with the lac Z coding region so that a hybrid - lac Z protein is produced; pIN vectors (Inouye & Inouye, *Nucleic Acids Res.*, 13:3101-3109, 1985; Van Heeke & Schuster, *J. Biol. Chem.* 264:5503-5509, 1989); glutathione-S-transferase (GST) and the like.

25 In yeast, a number of vectors containing constitutive or inducible promoters may be used. For a review see, *Current Protocols in Molecular Biology*, Vol. 2, 1988, Ed. Ausubel, et al., Green Publish. Assoc. & Wiley Interscience, Ch. 13; Grant, et al., 1987, *Expression and Secretion Vectors for Yeast*, in *Methods in Enzymology*, Eds. Wu & Grossman, 31987, Acad. Press, N.Y., Vol. 153, pp. 516-544; Glover, 1986,
30 DNA Cloning, Vol. II, IRL Press, Wash., D.C., Ch. 3; and Bitter, 1987, *Heterologous*

Gene Expression in Yeast, Methods in Enzymology, Eds. Berger & Kimmel, Acad. Press, N.Y., Vol. 152, pp. 673-684; and The Molecular Biology of the Yeast *Saccharomyces*, 1982, Eds. Strathern, et al., Cold Spring Harbor Press, Vols. I and II. A constitutive yeast promoter such as ADH or LEU2 or an inducible promoter such as GAL may be used (Cloning in Yeast, Ch. 3, R. Rothstein In: DNA Cloning Vol.11, A Practical Approach, Ed. DM Glover, 1986, IRL Press, Wash., D.C.). Alternatively, vectors may be used which promote integration of foreign DNA sequences into the yeast chromosome.

10 In cases where plant expression of the CACNA1G coding sequence may be driven by any of a number of promoters. For example, viral promoters such as the 35S RNA and 19S RNA promoters of CaMV (Brisson, et al., Nature 310:511-514, 1984), or the coat protein promoter to TMV (Takamatsu, et al., EMBO J.6:307-311, 1987) may be used; alternatively, plant promoters such as the small subunit of RUBISCO (Coruzzi, et al., EMBO J.3:1671-1680, 1984; Broglie, et al., Science 224:838-843, 15 1984); or heat shock promoters, e.g., soybean hsp17.5-E or hsp17.3-B (Gurley, et al., Mol. Cell. Biol. 6:559-565, 1986) may be used. These constructs can be introduced into plant cells using Ti plasmids, Ri plasmids, plant virus vectors, direct DNA transformation, microinjection, electroporation, etc. For reviews of such techniques see, for example, Weissbach & Weissbach, 1988, Methods for Plant Molecular 20 Biology, Academic Press, N.Y., Section VIII, pp. 421-463; and Grierson & Corey, 1988, Plant Molecular Biology, 2d Ed., Blackie, London, Ch. 7-9.

An alternative expression system which could be used to express is an insect 25 system. In one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The CACNA1G coding sequence may be cloned into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of the 30 CACNA1G coding sequence will result in inactivation of the polyhedrin gene and

production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted gene is expressed. (e.g., see Smith, et al., 1983, J. Virol. 46:584; Smith, U.S. Pat. No. 4,215,051).

5

Eukaryotic systems, and preferably mammalian expression systems, allow for proper post-translational modifications of expressed mammalian proteins to occur. Eukaryotic cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, phosphorylation, and advantageously, secretion of the gene product may be used as host cells for the expression of CACNA1G .
Mammalian cell lines may be preferable. Such host cell lines may include but are not limited to CHO, VERO, BHK, HeLa, COS, MDCK, -293, and W138.

Mammalian cell systems which utilize recombinant viruses or viral elements to direct expression may be engineered. For example, when using adenovirus expression vectors, the CACNA1G coding sequence may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the protein in infected hosts (e.g., see Logan & Shenk, Proc. Natl. Acad. Sci. USA, 81:3655-3659, 1984). Alternatively, the vaccinia virus 7.5K promoter may be used (e.g., see, Mackett, et al., 1982, Proc. Natl. Acad. Sci. USA 79:7415-7419; Mackett, et al., J. Virol. 49:857-864, 1984; Panicali, et al., Proc. Natl. Acad. Sci. USA 79:4927-4931, 1982). Of particular interest are vectors based on bovine papilloma virus which have the ability to replicate as extrachromosomal elements (Sarver, et al., Mol. Cell. Biol. 1:486, 1981). Shortly after entry of this DNA into mouse cells, the plasmid replicates to about 100 to 200 copies per cell. Transcription of the inserted cDNA does not require integration of the plasmid into the host's chromosome, thereby yielding a high level of expression. These vectors can be used for stable expression by

including a selectable marker in the plasmid, such as, for example, the neo gene. Alternatively, the retroviral genome can be modified for use as a vector capable of introducing and directing the expression of the CACNA1G gene in host cells (Cone & Mulligan, Proc. Natl. Acad. Sci. USA 81:6349-6353, 1984). High level expression
5 may also be achieved using inducible promoters, including, but not limited to, the metallothionine IIA promoter and heat shock promoters.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. Rather than using expression vectors which contain viral
10 origins of replication, host cells can be transformed with the CACNA1G cDNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and
15 grow to form foci which in turn can be cloned and expanded into cell lines. For example, following the introduction of foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, et al., Cell, 11:223, 1977), hypoxanthine-
20 guanine phosphoribosyltransferase (Szybalska & Szybalski, Proc. Natl. Acad. Sci. USA, 48:2026, 1962), and adenine phosphoribosyltransferase (Lowy, et al., Cell, 22: 817, 1980) genes can be employed in tk.sup.-, hgpert.sup.- or apert.sup.- cells respectively. Also, antimetabolite resistance can be used as the basis of selection for dhfr, which confers resistance to methotrexate (Wigler, et al., Natl. Acad. Sci. USA,
25 77:3567, 1980; O'Hare, et al., Proc. Natl. Acad. Sci. USA, 78: 1527, 1981); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, Proc. Natl. Acad. Sci. USA, 78: 2072, 1981; neo which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, et al., J. Mol. Biol., 150:1, 1981); and hygromycin (Santerre, et al., Gene, 30:147, 1984) genes. Recently,
30 additional selectable genes have been described, namely trpB, which allows cells to

utilize indole in place of tryptophan; hisD, which allows cells to utilize histinol in place of histidine (Hartman & Mulligan, Proc. Natl. Acad. Sci. USA, 85:8047, 1988); and ODC (ornithine decarboxylase) which confers resistance to the ornithine decarboxylase inhibitor, 2-(difluoromethyl)-DL-ornithine, DFMO (McConlogue L.,
5 1987, In: Current Communications in Molecular Biology, Cold Spring Harbor Laboratory, ed.).

Isolation and purification of microbial expressed polypeptide, or fragments thereof, provided by the invention, may be carried out by conventional means
10 including preparative chromatography and immunological separations involving monoclonal or polyclonal antibodies.

In one embodiment, the invention provides substantially purified polypeptide encoded by CACNA1G polynucleotide sequences. Exemplary CACNA1G
15 polypeptide is set forth in SEQ ID NO:52. The term "substantially purified" as used herein refers to a polypeptide which is substantially free of other proteins, lipids, carbohydrates or other materials with which it is naturally associated. One skilled in the art can purify a polypeptide encoded by CACNA1G polynucleotide sequence using standard techniques for protein purification. The substantially pure polypeptide
20 will yield a single major band on a non-reducing polyacrylamide gel. The purity of the CACNA1G polypeptide can also be determined by amino-terminal amino acid sequence analysis.

Minor modifications of the CACNA1G primary amino acid sequences may
25 result in proteins which have substantially equivalent activity as compared to the unmodified counterpart polypeptide described herein. Such modifications may be deliberate, as by site-directed mutagenesis, or may be spontaneous. All of the polypeptides produced by these modifications are included herein as long as the biological activity still ex-ists.

The polypeptides of the invention also include dominant negative forms of the CACNA1G polypeptide which do not have the biological activity of CACNA1G polypeptide sequence. A "dominant negative form" of CACNA1G is a polypeptide that is structurally similar to CACNA1G polypeptide but does not have wild-type CACNA1G function. For example, a dominant-negative CACNA1G polypeptide may interfere with wild-type CACNA1G function by binding to, or otherwise sequestering, regulating agents, such as upstream or downstream components, that normally interact functionally with the CACNA1G polypeptide.

10 Identification and Isolation of CACNA1G

To identify genes differentially methylated in colorectal cancer, methylated CpG island amplification was used followed by representational difference analysis (Razin and Cedar, Cell 17: 473-476, 1994, herein incorporated by reference). One of the clones recovered (MINT31, see U.S. Patent Application Serial No. 09/309,175, incorporated by reference herein in its entirety) mapped to human chromosome 17q21 using a radiation hybrid panel. A Blast search revealed this fragment to be completely identical to part of a BAC clone (Genbank: AC004590) sequenced by high throughput genomic sequencing. The region surrounding MINT31 fulfills the criteria of a CpG island: GC content 0.67, CpG/GpC ratio 0.78 and a total of 305 CpG sites in a 4 kb region. Using this CpG island and 10 kb of flanking sequences in a Blast analysis, several regions highly homologous to the rat T-type calcium channel gene, CACNA1G, were identified (Perez-Reyes et al., Nature 391: 896-900, 1998, herein incorporated by reference). Several ESTs were also identified in this region. Using Genscan, 2 putative coding sequences (G1, and G2) were identified. Blastp analysis revealed that G1 has a high homology to the EH-domain-binding protein, epsin, while G2 is homologous to a C. elegans hypothetical protein (accession No. 2496828).

The MINT31 CpG island corresponds to the 3' regions of G1 and G2, based on the direction of the open reading frame and the presence of a poly A tail, and, without

being bound by theory, is unlikely to influence their transcription. The EST closest to MINT31 (H13333) (SEQ ID NO:120; Figure 5) was sequenced entirely and was found not to contain a continuous open reading frame, but a poly-adenylation signal was identified on one end, along with a poly A tail. These data suggest that H13333 corresponds to the last 2 exons of an unidentified gene. MINT31 is in the intron of this gene and is, again without being bound by theory, unlikely to influence transcription of the unidentified gene. However, based on both promoter prediction (TSSG) analysis of this region and homology to the rat CACNA1G sequence, the MINT31 CpG island is also in the 5' region of human CACNA1G gene and is likely to play a role in its transcriptional activity.

The human CACNA1G sequence deposited in Genbank lacks the 5' region of the gene, when compared to the rat homologue. To determine the 5' region of human CACNA1G, cDNA was amplified by RT-PCR using primers based on the BAC sequence (Genbank: AC004590, herein incorporated by reference). The PCR products were cloned and sequenced, and the genomic organization of the gene was determined by comparing the newly identified sequences as well as the known sequences to the BAC that covers this region. CACNA1G is composed of 34 exons which span a 70 kb area. (See, Figure 3A and SEQ ID NO:52). Based on sequences deposited in Genbank, the gene has two possible 3' ends caused by alternate splicing. Human CACNA1G is highly homologous to rat CACNA1G with 93% identity at the protein level, and 89% identity at the nucleotide level. The 5' flanking region of human CACNA1G lacks TATA and CAAT boxes, which is similar to many housekeeping genes. A putative TFIID binding site was identified 547-556 bp upstream from the translation start site, and several other potential transcription factor binding sites such as AP1 (1 site), AP2 (2 sites) and SP1 (10 sites), were identified upstream of CACNA1G exon 1 using the promoter prediction program, TESS.

Methylation Analysis of CACNA1G

The CACNA1G CpG island is 4 kb, and is larger than many typical CpG islands. MINT31 corresponds to the 5' edge of the island while CACNA1G is in the 3' region. It is not known whether large CpG islands such as this are coordinately regulated with regards to protection from methylation, and aberrant methylation in cancer.

To address this issue, the methylation status of the 5' region of CACNA1G was studied using bisulfite-PCR of DNA from normal tissues as well as 35 human cancer cell lines from colon, lung, prostate, breast and hematopoietic tumors. More specifically, forty-nine primary colorectal cancers, 28 colorectal adenomas, 16 primary gastric cancers and 17 acute myelogenous leukemia samples were used for methylation analyses. DNA from eight colon cancer cell lines (Caco2, RKO, SW48, HCT116, DLD1, Lovo, SW837, HT29), 4 lung cancer cell lines (OH3, H249, H157, H209), 4 glioblastoma cell lines (Daoy, D283, U87, U373), 8 breast cancer cell lines (MB-468, MCF7, MB-231, MB-474, MB-435, MB-453, BT20, CAMA1, SKBR3), 7 hematopoietic tumor cell lines (CEM, Raji, KG1A, HL60, ML-1, Molt3, K562), and 4 prostate cancer cell lines (DU145, DUPRO, LNCAP, TSUPRL) were also investigated. The CpG island was divided into 8 regions (SEQ ID NOs: 35-42, respectively). The methylation status of each region was examined separately. The genomic DNA was treated with sodium bisulfite and PCR amplified using primers containing no or a minimum number of CpG sites. (For a detailed description of bisulfite-PCR, see, U.S. Patent No. 5,786,146, incorporated herein by reference in its entirety). Methylated alleles were detected by digesting the PCR products using restriction enzymes which specifically cleave sites created or retained due to the presence of methylated CpGs. None of the regions was methylated in normal colon, consistent with a uniform protection against de-novo methylation.

Regions 1 and 2 were frequently methylated in cancer cell lines, and behaved in a concordant manner with respect to methylation pattern. Indeed, these 2 regions were methylated in most cancer cell types except gliomas. Moreover, most cell lines where methylation was found methylated both regions 1 and 2. in contrast, region 3, which is less CG rich than any of the other regions, had either no methylation or very low levels of methylation in most cell lines. Regions 5, 6, and 7 behaved quite differently compared to 1-3. Methylation of these regions was less frequent than regions 1-2, as 22/35 cell lines had no detectable methylation there, despite often showing methylation of region 1-2. However, when methylation of regions 5, 6, or 7 was present (in 13/35 cell lines), it affected all 3 regions in a coordinate manner, although to varying extents. Finally, regions 4 and 8 behaved differently again, being partially methylated primarily in colon and breast cell lines. Therefore, with regards to hypermethylation in cancer cells, the CpG rich region upstream of CACNA1G appears to be composed of 2 CpG islands which behave independently. MINT31 corresponds to the upstream CpG island (island 1, regions 1 and 2), while the 5' region of CACNA1G is contained in the downstream CpG island (island 2, regions 5-7). Regions 3, 4 and 8 correspond to the edge of these CpG islands, and behave a little differently than the hearts of the CpG islands, as previously described for the E-Cad gene (Graff, et al., J. Biol. Chem. 272: 22322-22329, 1997).

20

Overall, the methylation patterns of CACNA1G fell into 5 distinct categories: (1) No methylation in any region (normal tissue). (2) Slight methylation of island 1 (6 cell lines, e.g., TSU-PRL). (3) Heavy methylation of island 1 but no methylation of island 2 (16 cell lines, e.g., Caco2). (4) Heavy methylation of island 1 and moderate to heavy methylation of island 2 (6 cell lines, e.g., RKO and Raji). (5) High methylation of island 1 and low to moderate methylation of island 2 (7 cell lines, e.g., MB-231).

30

Methylation Dependent Expression of CACNA1G

In a previous study, rat CACNA1G was shown to be expressed most abundantly in the brain (Perez-Reyes et al., Nature 391: 896-900. 1998). To determine the expression of CACNA1G in normal and neoplastic human cells, RT-PCR was performed using cDNA from various normal tissues and from a panel of 27 tumor cell lines. CACNA1G was expressed ubiquitously in a variety of tissues and cell lines. In normal tissues expression was relatively low but easily detectable, while most cell lines had relatively high expression of CACNA1G. However, some cell lines had negligible or totally absent levels of CACNA1G expression. The results of CACNA1G expression was correlated with the detailed methylation analysis previously described. In this analysis, a remarkable pattern emerged. Methylation of regions 1-4 and 8 had no effect on CACNA1G expression. However, there was a strong correlation between methylation of regions 5-7 and expression of the gene. In fact, all cell lines tested that lack methylation of this region strongly express the gene. All 6 cell lines with pattern 4 methylation studied had no detectable expression. Finally, the 7 cell lines with pattern 5 methylation (examples DLD-1 and MB-453) had variable levels of expression ranging from very low to near normal. The fact that patterns 3 and 5 differ significantly with regards to expression, but are almost identical with regards to methylation of all regions except 7 indicates that this area is important in the inactivation of CACNA1G.

To confirm that methylation of the 5' CpG island of CACNA1G is associated with gene inactivation, 3 non-expressing cell lines showing pattern 4 methylation (RKO, SW48 and Raji) and 2 weakly expressing cell lines showing pattern 5 methylation (MB-231 and MB-435) were treated with 1 M of the methyl-transferase inhibitor 5-deoxy-azacitidine. After treatment, all of these cell lines re-expressed CACNA1G mRNA. Consistent with re-expression, demethylation of region 7 was observed after 5-deoxy-azacitidine treatment.

De novo cytosine methylation is thought to sometimes occur in vitro during cell propagation (Antequera et al., Cell 62: 503-514, 1990). To determine whether the methylation of CACNA1G occurs in vivo, primary human tumors were examined for methylation of the 5' region of CACNA1G. Aberrant methylation was detected in 17
5 out of 49 (35%) colorectal cancers, 4 out of 28 colorectal adenomas (25%), 4 out of 16 (25%) gastric cancers and 3 out of 17 (18%) acute myelogenous leukemia cases. In colorectal cancers, there was a significant correlation between methylation of CACNA1G and methylation of p16 ($p < 0.005$) and hMLH1 ($p < 0.001$), as well as a strong correlation with the presence of microsatellite instability, and the recently
10 identified CpG island methylator phenotype (CIMP), indicating that CACNA1G is also a target for CIMP in colorectal cancer.

To determine whether aberrant methylation of the 5' region of CACNA1G affects the expression status of this gene in primary tumors, RT-PCR was performed
15 using cDNA from a series of colorectal adenomas. Six out of 8 cases which showed no methylation of region 7 expressed CACNA1G. In sharp contrast, all 5 cases that showed methylation of region 7 had no detectable expression of the gene.

Thus, a human T-type calcium channel gene (CACNA1G) has been identified
20 and cloned using the MINT31 sequence as a probe. The human T-type calcium channel gene has been determined to be a target of aberrant methylation and silencing in human tumors. The data show that MINT31 (for a representative sequence of MINT1-33, see, US Patent Application Serial No. 09/309,175) hereby incorporated by reference can be used as a probe to identify genes that play a role in disorders such as
25 cell proliferative disorders.

Detailed analysis of the CpG island upstream of CACNA1G revealed that methylation 300 to 800 bp upstream of the gene closely correlated with transcriptional inactivation. The CACNA1G promoter is contained in a large CG rich area that is not
30 coordinately methylated in cancer. The CpG island around MINT31 is much more

frequently methylated in cancers compared to that just upstream of CACNA1G. This may simply be caused by differential susceptibility to de-novo methylation between these two regions, with methylation of MINT31 serving as a trigger, and eventually spreading to CACNA1G, as described in other genes (Graff, et al., J. Biol. Chem. 272: 22322-22329, 1997). However, it is likely that these 2 regions are controlled by different mechanisms because (1) cell lines kept in culture for countless generations do not in fact spread methylation from MINT31 to CACNA1G (e.g., Caco2), (2) region 3 that separates the 2 islands is infrequently and sparsely methylated in cancer and (3) 2 cases of primary colorectal cancer were found which are methylated at the CACNA1G promoter but not at MINT31). Therefore, methylation of MINT31 appears to be independent of methylation of CACNA1G suggesting that they are 2 distinct CpG islands regulated by different mechanisms.

Many CpG islands of silenced genes appear to be methylated uniformly and heavily throughout the island (e.g., Graff, et al., J. Biol. Chem. 272: 22322-22329, 1997). In contrast the methylation patterns of the 5' region of CACNA1G (region 5-7) was heterogeneous in the cell lines which did not express this gene. Nevertheless, methylation clearly plays a role in CACNA1G repression since demethylation readily reactivates the gene.

20

The mechanism of CACNA1G methylation remains to be determined. Methylation was not detected in normal colon mucosa, placenta, normal breast epithelium and normal bone marrow, including samples from aged patients, suggesting that methylation of this region is specific for cell proliferative disorders such as cancer, and the like. However, there was a significant correlation between methylation of CACNA1G and other tumor suppressor genes such as p16 and hMLH1. Thus, CACNA1G is likely a target for the recently described CIMP phenotype, which results in a form of epigenetic instability with simultaneous inactivation of multiple genes.

30

T-type calcium channels are involved not only in electrophysiological rhythm generation but also in the control of cytosolic calcium during cell proliferation and cell death (reviewed in Berridge, et al., *Nature* 395: 645-648, 1998). Expression of CACNA1G is not limited to brain and heart, indicating a likely role in other tissues in which it is expressed. It has previously been shown that Ca^{2+} influx via T-type channels is an important factor during the initial stages of cell death such as apoptosis (Berridge, et al., *Nature* 395: 645-648, 1998), ischemia (Fern, *J. Neurosci.* 18: 7232-7243, 1998) and complement-induced cytotoxicity (Newsholme, et al., *Biochem. J.* 295: 773-779, 1993.). The studies culminating in the present invention indicates that impairment of voltage gated calcium channels plays an important role in cancer development and progression through altering calcium signaling.

Due to the clear correlation between methylation of CpG islands and cellular proliferative disorders, in another embodiment of the present invention, there are provided methods for detecting a cellular proliferative disorder in a subject having or at risk for said cellular proliferative disorder. The method includes assaying, in nucleic acid-containing specimen taken from said subject, the methylation state of a gene or its associated regulatory regions, wherein the expression state of the gene or its associated regulatory regions is associated with the presence of the cellular proliferative disorder, and identifying as having a cellular proliferative disorder a subject that has aberrant methylation of regions of said gene. The method provides for detecting a cellular proliferative disorder in a subject having or at risk for said cellular proliferative disorder by identifying aberrantly methylation of regions of a gene when compared to the same regions of the gene in a subject not having said cellular proliferative disorder.

The aberrant methylation comprises hypermethylated CpG rich regions (i.e., islands). In one aspect of the present invention, the CpG rich regions are associated with the CACNA1G gene, and affect the expression thereof in a methylation state-dependent manner. A "cell proliferative disorder" or "cellular proliferative disorder"

is any disorder in which the proliferative capabilities of the affected cells is different from the normal proliferative capabilities of unaffected cells. An example of a cell proliferative disorder is neoplasia. Malignant cells (i.e., cancer) develop as a result of a multistep process. Specific, non-limiting examples of cell proliferative disorders associated with increased methylation of CpG-islands are low grade astrocytoma, anaplastic astrocytoma, glioblastoma, medulloblastoma, gastric cancer, colorectal cancer, colorectal adenoma, acute myelogenous leukemia, lung cancer, renal cancer, leukemia, breast cancer, prostate cancer, endometrial cancer and neuroblastoma.

10 A cell proliferative disorder as described herein may be a neoplasm. Such neoplasms are either benign or malignant. The term "neoplasm" refers to a new, abnormal growth of cells or a growth of abnormal cells that reproduce faster than normal. A neoplasm creates an unstructured mass (a tumor) which can be either benign or malignant. For example, the neoplasm may be a head, neck, lung, esophageal, stomach, prostate, small bowel, colon, bladder, kidney, or cervical
15 neoplasm. The term "benign" refers to a tumor that is noncancerous, e.g. its cells do not proliferate or invade surrounding tissues. The term "malignant" refers to a tumor that is metastatic or no longer under normal cellular growth control.

20 A cell proliferative disorder may be an age-associated disorder. Examples of age-associated disorders which are cell proliferative disorders include colon cancer, lung cancer, breast cancer, prostate cancer, leukemia and melanoma, amongst others.

A "nucleic acid containing specimen" includes any type of material containing
25 a nucleic acid to be subject to invention methods. The nucleic acid may be contained in a biological sample. Such samples include but are not limited to any bodily fluid, such as a serum, urine, saliva, blood, cerebrospinal fluid, pleural fluid, ascites fluid, sputum, stool, or a biopsy sample.

30 Samples or specimens include any CpG-rich DNA sequence, whatever the origin, as long as the sequence is detectably present in a sample. While routine

diagnostic tests may not be able to identify cancer cells in these samples, the method of the present invention identifies neoplastic cells derived from the primary tumor or from a metastases. The method includes non-invasive sampling (e.g., bodily fluid) as well as invasive sampling (e.g., biopsy). The sample of DNA of the subject may be
5 serum, plasma, lymphocytes, urine, sputum, bile, stool, cervical tissue, saliva, tears, cerebral spinal fluid, regional lymph node, histopathologic margins, and any bodily fluid that drains a body cavity or organ. Therefore, the method provides for the non-invasive detection of various tumor types including head and neck cancer, lung cancer, esophageal cancer, stomach cancer, small bowel cancer, colon cancer, bladder
10 cancer, kidney cancers, cervical cancer and any other organ type that has a draining fluid accessible to analysis. For example, neoplasia of regional lymph nodes associated with a primary mammary tumor can be detected using the method of the invention. Regional lymph nodes for head and neck carcinomas include cervical lymph nodes, prelaryngeal lymph nodes, pulmonary juxta-esophageal lymph nodes
15 and submandibular lymph nodes. Regional lymph nodes for mammary tissue carcinomas include the axillary and intercostal nodes. Samples also include urine DNA for bladder cancer or plasma or saliva DNA for head and neck cancer patients.

Any nucleic acid sample, in purified or nonpurified form, can be utilized as the
20 starting nucleic acid or acids in accordance with the present invention, provided it contains, or is suspected of containing, a nucleic acid sequence containing a target locus (e.g., CpG-containing nucleic acid). In general, the CpG-containing nucleic acid is DNA. However, invention methods may employ, for example, samples that contain DNA, or DNA and RNA, including messenger RNA, wherein DNA or RNA
25 may be single stranded or double stranded, or a DNA-RNA hybrid may be included in the sample. A mixture of nucleic acids may also be employed. The specific nucleic acid sequence to be detected may be a fraction of a larger molecule or can be present initially as a discrete molecule, so that the specific sequence constitutes the entire nucleic acid. It is not necessary that the sequence to be studied be present initially in a
30 pure form; the nucleic acid may be a minor fraction of a complex mixture, such as

contained in whole human DNA. The nucleic acid-containing sample used for detection of methylated CpG may be from any source including, but not limited to, brain, colon, urogenital, lung, renal, pancreas, liver, esophagus, stomach, hematopoietic, breast, thymus, testis, ovarian, prostate and uterine tissue, and may be
5 extracted by a variety of techniques such as that described by Maniatis, et al. (Molecular Cloning: a Laboratory Manual, Cold Spring Harbor, NY, pp 280, 281, 1982).

The nucleic acid of interest can be any nucleic acid where it is desirable to
10 detect the presence of a differentially methylated CpG island. The CpG island comprises a CpG island located in a gene or regulatory region for a gene. A "CpG island" is a CpG rich region of a nucleic acid sequence. The nucleic acid sequence may include, for example, APOB, CACNA1G, CDX2, EGFR, FBN1, GPR37, HSPA6, IQGAP2, KL, PAR2, PITX2, PTCH, SDC1, or SDC4 (see for example
15 Figures 4A-4F). Alternatively the nucleic acid of interest can be, for example, a MINT31 nucleic acid sequence (SEQ ID NO:120. However, any gene or nucleic acid sequence of interest containing a CpG sequence can provide diagnostic information (i.e., via its methylation state) using invention methods.

Moreover, these markers can also be multiplexed in a single amplification
20 reaction to generate a low cost, reliable cancer screening test for many cancers simultaneously. A combination of DNA markers for CpG-rich regions of nucleic acid may be amplified in a single amplification reaction. The markers are multiplexed in a single amplification reaction, for example, by combining primers for more than one
25 locus. For example, DNA from a urine sample can be amplified with three different randomly labeled primer sets, such as those used for the amplification of the CACNA1G, EGFR and PTCH loci, in the same amplification reaction. The reaction products are separated on a denaturing polyacrylamide gel, for example, and then exposed to film for visualization and analysis. By analyzing a panel of markers, there
30 is a greater probability of producing a more useful methylation profile for a subject.

If the sample is impure (e.g., plasma, serum, stool, ejaculate, sputum, saliva, cerebrospinal fluid, or blood or a sample embedded in paraffin), it may be treated before amplification with a reagent effective for lysing the cells contained in the fluids, tissues, or animal cell membranes of the sample, and for exposing the nucleic acid(s) contained therein. Methods for purifying or partially purifying nucleic acid from a sample are well known in the art (e.g., Sambrook et al., Molecular Cloning: a Laboratory Manual, Cold Spring Harbor Press, 1989, herein incorporated by reference).

10 In order to detect a differential methylation state for a gene or CpG-containing region of interest, invention methods include any means known in the art for detecting such differential methylation. For example, detecting the differential methylation may include contacting the nucleic acid-containing specimen with an agent that modifies unmethylated cytosine, amplifying a CpG-containing nucleic acid in the specimen by
15 means of CpG-specific oligonucleotide primers, wherein the oligonucleotide primers distinguish between modified methylated and nonmethylated nucleic acid, and detecting the methylated nucleic acid based on the presence or absence of amplification products produced in said amplifying step. This embodiment includes the PCR-based methods described in U.S. Patent No. 5,786,146, incorporated herein
20 in its entirety.

For the first time, the methylation state of a number of genes has been correlated with cell proliferative disorders. Examples of such genes, primers useful for identifying their methylation state, and general PCR conditions are set out in Table
25 1.

TABLE 1 Bisulfite-PCR Primers

Genes	5'-primer	3'-primer	Annealing Temperature(°C)
APOB	5'-gtttttagttaggggtttt-3'	5'-RccaacaacRccaacaac-3'	57
	TaqI		
5 CACNA1G	See Table 2		
R6	5'-tgttgggttgggatagtt-3'	5'-acraaaaaaaaaaaaaaatctt-3'	47TD
	HinfI		
R7	5'-ggggggttttttyggattt-3'	5'-ttccctacrcctctaaactcc-3'	49TD
	EcoRI		
10 CDX2	5'-gtaggttagaggaggatyggt-3'	5'-aaaacaaacctcaccatactac-3'	60
	TaqI		
EGFR	5'-tttgattYgttagtattgat-3'	5'-cccttaccttcttttct-3'	52
	HinfI, TaqI		
FBN1	5'-ttttttattgYgttaatttg-3'	5'-ttccctcacctcttcaata-3'	54
15	TaqI		
GPR37	5'-ggtaggtgggtaagagttt-3'	5'-aacRtttaattcaatacaaac-3'	56
	HinfI		
HSPA6	5'- tttttagtagtttgagttagagg-3'	5'- ttctatccttttacacctccc-3'	60
	TaqI		
20 IQGAP2	5'-tttattttatttagtgtag-3'	5'-ctcttcRtataacatcctac-3'	52
	TaqI		
KL	5'- gtagtatgttygttagygtt-3'	5'- aaataccctaaaaaaaccc-3'	60
	TaqI		
PAR2	5'-YggttttttGaaatttaattc-3'	5'-aactccRcatcctctctaaa-3'	45
25	HinfI, TaqI		
PITX2	5'- taagtgtttagtaggtttt-3'	5'- ccaaactccactacacaataac-3'	60
	TaqI		
PTCHA	5'- gttggttgttaatyggagt-3'	5'- ttaccaaccraaacatatt-3'	60
	TaqI		
30 PTCHB	5'- aatgtgtggaatttagga-3'	5'- taaaacaaccaactacaacttac-3'	60
	TaqI		
SDC1	5'- agaggaattyggttagtagag-3'	5'- cacaacaaaacaaaac-3'	60
	TaqI		
SDC4	5'-ggggattgttggtagtgg-3'	5'-cccgaattccccaataaa-3'	56
35	HinfI		

Note: Y = C or T, R = G or A; row 1-SEQ ID NO:1 and 2; row 2-SEQ ID NO:3 and 4 and so forth through to SEQ ID NO:32, respectively. The gene sequences can be found in a gene database found at <http://www.ncbi.nlm.nih.gov/UniGene/index.html>.

40

In one embodiment, the oligonucleotide primers are specifically targeted to CACNA1G and its associated CpG islands as described herein. Examples of oligonucleotides suited for determining the methylation state of the 8 regions of the two CpG islands of MINT31/CACNA1G, as well as PCR conditions and useful
45 methylation sensitive restriction endonucleases are set out in Table 2.

TABLE 2 Primers Useful For Bisulfite/PCR Analysis of CACNA1G

Region	Primer set, forward /reverse	Annealing temperature (cycles)	Restriction enzyme
5	Region 1 5'-GAYGGYGTAGTAGTTATTTTGTT-3'	58 (3), 56 (4),	BstUI
	F165 5'-CATCACCACCCCTCACTTTAC-3'	54 (5), 52 (26)	MaeII
	Region 2 5'-TTYGGGTATTTATAGTTTTTGGAG-3'	60 (3), 58 (4),	TaqI
	GM2 5'-AATCTACRCCTTCACTCACTC-3'	56 (5), 54 (26)	BstUI
	Region 3 5'-TTTAGGAGYGTTAATGTGAGGTT-3'	55 (3), 53 (4),	HinfI
10	GM3 5'-CTAAAAAAACCCAATCTTAAAAAAAC-3'	51(5), 49(26)	MaeII
	Region 4 5'-TGGATAAAGGATGTTTGGGGTTTG-3'	55 (5), 53 (5),	MaeII
	GM5 5'-CCCTCCCCTTACCCCTAAATCC-3'	51(5), 49(26)	TaqI
	Region 5 5'-AATYGGATTTTAGTTGTGGTTTTT-3'	60 (3), 58 (4),	BstUI
	GM1 5'-CACACCACAATAAATCCCTCACT-3'	56 (5), 54 (26)	TaqI
15	Region 6 5'-TTGTGGYGTTGGYGATAGTT-3'	53 (3), 51 (4),	HinfI
	GM6 5'-ACRAAAAAAAAAAAAAAAAAATCTCTT-3'	49 (5), 47 (26)	
	Region 7 5'-GGGGGYGTTTTTTTTYGGATTTT-3'	55 (5), 53 (5),	EcoRI
	GM4 5'-TTCCCCTACRCCCCTAAACTTCC-3'	51(5), 49(26)	
	Region 8 5'-GGGAGTTTGGGAGTTGTATTTTGTT-3'	60 (3), 58 (4),	TaqI
20	5'-AACCAAATTAAAAAATCAAACCCTAA-3'	56 (5), 54 (26)	
	Region 8 5'-GAGGGGGGATYGTAATTTTTTG-3'	60 (3), 58 (4),	BstUI
	5'-CCRAATCTCCTTATTATACCTCCAA-3'	56 (5), 54 (26)	

25 Region 1 F/R=SEQ ID NO:33 and 34; Region 2=SEQ ID NO:35 and 36 and so forth through
Region 8 (TaqI)=SEQ ID NO:47 and 48 and Region 8 (BstUI)=SEQ ID NO:49 and 50.

Exemplary target regions (i.e., regions 1-8 of MINT31/CACNA1G) that are complementary to the primers listed in Table 2 are provided in Table 3:

Table 3 - Target Sequences

Region/ Primer	Target set, forward /reverse	SEQ ID NO	Corresponding Primer SEQ ID NO
Region 1	5'- AACAAAATAACTACTACRCCRTC-3'	87	33
F165	5'- GTAAAGTGAGGGGTGGTGATG-3'	88	34
Region 2	5'- CTCCAAAAAATACTATAAATACCCRAA-3'	89	35
GM2	5'- GAGTGAGTGAAGGYGGTAGATT-3'	90	36
Region 3	5'- AACCTCACATTAACRCTCCTAAA-3'	91	37
GM3	5'- GTTTTTTTAAGATTGGGTTTTTTTAG-3'	92	38
Region 4	5'- CAAACCCCAAACATCCTTTATCCA-3'	93	39
GM5	5'- GGATTTAGGGGTAAGGGGAGGG-3'	94	40
Region 5	5'- AAAAACCACAATAAAATCCRATT-3'	95	41
GM1	5'- AGTGAGGGATTTAGTTGTGGTGTG-3'	96	42
Region 6	5'- AACTATCRCCAACRCCACAA-3'	97	43
GM6	5'- AAGAGATTTTTTTTTTTTTTTTTTYGT-3'	98	44
Region 7	5'- AAAATCCRAAAAAAAAAACRCCCCC-3'	99	45
GM4	5'- GGAAGTTTAGGGGYGTAGGGGAA-3'	100	46
Region 8	5'- AACAAAATACAACCTCCAAACACCC-3'	101	47
	5'- TTAGGGTTTGATTTTTTAATTTGGTT-3'	102	48
Region 8	5'- CAAAAAATTACRATCCCCCCTC-3'	103	49
	5'- TTGGAGGTATAATAAGGAGATTTTGG-3'	104	50

TABLE 4 Targets for Bisulfite-PCR Primers

Genes		SEQ ID NO	Corresponding Primer SEQ ID NO
APOB 5'-target	5'-AAAAAACCCAAACTACAAAAAC-3'	55	1
3'-target	5'GTTGTTGGRGTTGTTGGR -3'	56	2
R6 5'-target	5'- AACTATCYCCAACYCCACAA -3'	57	3
3'-target	5'- AAGAGATTTTTTTTTTTTTTTTTRGT -3'	58	4
R7 5'-target	5'- AAAATCCYAAAAAAAACYCCCCC -3'	59	5
3'-target	5'- GGAAGTTTTAGGGGRGTAGGGGAA -3'	60	6
CDX2 5'-target	5'- AACYATCCCTCCCTCTAACCTAC -3'	61	7
3'-target	5'- AGGTAGTATGGTGAGGTTTGTTTT -3'	62	8
EGFR 5'-target	5'- ATCAATACTAAACRAAATCAAA -3'	63	9
3'-target	5'- AGGAAAAGAAAGGTAAGGG -3'	64	10
FBN1 5'-target	5'- CAAAATTAACRCAATAAAAAAA -3'	65	11
3'-target	5'- TATTTGAAGAGGTGGGAAA -3'	66	12
GPR37 5'-target	5'- AAACCTTACCCACCTAACC -3'	67	13
3'-target	5'- GGTGTGTAATTGGATTAAAYGTT -3'	68	14
HSPA6 5'-target	5'- CCACTAACTCAAAACTAAAAAA -3'	69	15
3'-target	5'- GGGAGGTGTAAAAGGTGAAA -3'	70	16
IQGAP2 5'-target	5'- CTAACACTAAAATAAAAAATAAA -3'	71	17
3'-target	5'- GTAGGATGTTATAYGAAGAG -3'	72	18
KL 5'-target	5'- AAACRCTAACRAACATACTAC -3'	73	19
3'-target	5'- GGGTTTTTTTTAGGGTATTT -3'	74	20
PAR2 5'-target	5'- GAATTAAATTTCAAAAACCR-3'	75	21
3'-target	5'- TTTAGGAGGATGYGGAGTT -3'	76	22
PITX2 5'-target	5'- AAAAAACCTAACRAAACACTTA -3'	77	23
3'-target	5'- GTTATTGTGTAGTGGAGTTTGG -3'	78	24
PTCHA 5'-target	5'- ACTCCRATTAACAAACCAAC -3'	79	25
3'-target	5'- AATATGGTTTYGGTTGGTAA -3'	80	26
PTCHB 5'-target	5'- TCCCTAAATCCACACATT -3'	81	27
3'-target	5'- GTAAGTTGTAGTTGGTTGTTTA -3'	82	28
SDC1 5'-target	5'- CTCTCTACTACCRAATTCCTCT -3'	83	29
3'-target	5'- GTTTTGGTTTGGTTGTG -3'	84	30
SDC4 5'-target	5'- CCACTACCAACAAATCCCC -3'	85	31
3'-target	5'- TTTATTGGGGAATTTCCGGG -3'	86	32

Table 5 New genes differentially methylated in disease versus normal issue

Gene Symbol	Gene name	Map	Unigene Entry ¹	Methylated In ²
<i>APOB</i>	Apolipoprotein B	2p24	Hs.585	Common Tumors
<i>CACNAIG</i>	T-type calcium channel	17	-	
<i>CDX2</i>	Caudal type homeo box transcription factor 2	13q12,3	Hs.77399	Leukemias, breast, prostate
<i>EGFR</i>	Epidermal Growth Factor Receptor	7p12	Hs.77432	Leukemias, breast
<i>FBN1</i>	Fibrillin-1	15q21.1	Hs.750	Colon, Breast, prostate, leukemias
<i>GPR37</i>	G protein-coupled receptor 37	7q31	Hs.27747	colon, breast, leukemias
<i>HSPA6</i>	Heat shock 70kD protein 6 (HSP70B')	1q	Hs.3268	Common tumors
<i>IQGAP2</i>	RasGAP-related protein	5q	Hs.78993	Common tumors
<i>KL</i>	Klotho	-	Hs.94592	Common tumors
<i>PAR2</i>	Proteinase-activated receptor 2	5q13	Hs.15429 9	Leukemias, breast
<i>PITX2</i>	Paired-like homeodomain transcription factor 2	4q25-27	Hs.92282	Leukemias, prostate, breast
<i>PTCH³</i>	Patched	9Q31	Hs.15952 6	Leukemias
<i>SDC1</i>	Syndecan 1	2p24.1	Hs.82109	Leukemias
<i>SDC4</i>	Syndecan 4	20q12	Hs.72082	Leukemias

Table 5: New genes differentially methylated in cancer and other diseases.

¹ Gene database that can be found at <http://www.ncbi.nlm.nih.gov/UniGene/index.html>.² Examples: List is not comprehensive.³ Two promoters are affected.

In another embodiment, detection of differential methylation is accomplished by contacting a nucleic acid sample suspected of comprising a CpG-containing nucleic acid with a methylation sensitive restriction endonuclease that cleaves only unmethylated CpG sites under conditions and for a time to allow cleavage of unmethylated nucleic acid. The sample is further contacted with an isoschizomer of the methylation sensitive restriction endonuclease, that cleaves both methylated and unmethylated CpG-sites, under conditions and for a time to allow cleavage of methylated nucleic acid. Oligonucleotides are added to the nucleic acid sample under conditions and for a time to allow ligation of the oligonucleotides to nucleic acid cleaved by the restriction endonuclease, and the digested nucleic acid is amplified by conventional methods such as PCR wherein primers complementary to the oligonucleotides are employed. Following identification, the methylated CpG-containing nucleic acid can be cloned, using method well known to one of skill in the art (see Sambrook et al., Molecular Cloning: a Laboratory Manual, Cold Spring Harbor Press, 1989).

As used herein, a "methylation sensitive restriction endonuclease" is a restriction endonuclease that includes CG as part of its recognition site and has altered activity when the C is methylated as compared to when the C is not methylated. Preferably, the methylation sensitive restriction endonuclease has inhibited activity when the C is methylated (e.g., SmaI). Specific non-limiting examples of a methylation sensitive restriction endonucleases include Sma I, BssHII, or HpaII. Such enzymes can be used alone or in combination. Other methylation sensitive restriction endonucleases will be known to those of skill in the art and include, but are not limited to SacII, EagI, and BstUI, for example. An "isoschizomer" of a methylation sensitive restriction endonuclease is a restriction endonuclease which recognizes the same recognition site as a methylation sensitive restriction endonuclease but which cleaves both methylated and unmethylated CGs. One of skill in the art can readily determine appropriate conditions for a restriction endonuclease to cleave a nucleic acid (see Sambrook et al., Molecular Cloning: a Laboratory Manual, Cold Spring

Harbor Press, 1989). Without being bound by theory, actively transcribed genes generally contain fewer methylated CGs than in other genes.

In one embodiment of the invention, a nucleic acid of interest is cleaved with a
5 methylation sensitive endonuclease. In one aspect, cleavage with the methylation
sensitive endonuclease creates a sufficient overhang on the nucleic acid of interest.
Following cleavage with the isoschizomer, the cleavage product can still have a
sufficient overhang. An "overhang" refers to nucleic acid having two strands wherein
the strands end in such a manner that a few bases of one strand are not base paired to
10 the other strand. A "sufficient overhang" refers to an overhang of sufficient length to
allow specific hybridization of an oligonucleotide of interest. In one embodiment, a
sufficient overhang is at least two bases in length. In another embodiment, the
sufficient overhang is four or more bases in length. An overhang of a specific
sequence on the nucleic acid of interest may be desired in order for an oligonucleotide
15 of interest to hybridize. In this case, the isoschizomer can be used to create the
overhang having the desired sequence on the nucleic acid of interest.

In another aspect of this embodiment, the cleavage with a methylation
sensitive endonuclease results in a reaction product of the nucleic acid of interest that
20 has a blunt end or an insufficient overhang. In this embodiment, an isoschizomer of
the methylation sensitive restriction endonuclease can create a sufficient overhang on
the nucleic acid of interest. "Blunt ends" refers to a flush ending of two stands, the
sense stand and the antisense strand, of a nucleic acid.

25 Once a sufficient overhang is created on the nucleic acid of interest, an
oligonucleotide is ligated to the nucleic acid cleaved of interest which has been
cleaved by the methylation specific restriction endonuclease. "Ligation" is the
attachment of two nucleic acid sequences by base pairing of substantially
complementary sequences and/or by the formation of covalent bonds between two
30 nucleic acid sequences. In one aspect of the present invention, an "oligonucleotide" is

a nucleic acid sequence of about 2 up to about 40 bases in length. It is presently preferred that the oligonucleotide is from about 15 to 35 bases in length.

In one embodiment, an adaptor is utilized to create DNA ends of desired sequence and overhang. An "adaptor" is a double-stranded nucleic acid sequence with one end that has a sufficient single-stranded overhang at one or both ends such that the adaptor can be ligated by base-pairing to a sufficient overhang on a nucleic acid of interest that has been cleaved by a methylation sensitive restriction enzyme or an isoschizomer of a methylation sensitive restriction enzyme. Adaptors can be obtained commercially, or two oligonucleotides can be utilized to form an adaptor. Thus, in one embodiment, two oligonucleotides are used to form an adaptor; these oligonucleotides are substantially complementary over their entire sequence except for the region(s) at the 5' and/or 3' ends that will form a single stranded overhang. The single stranded overhang is complementary to an overhang on the nucleic acid cleaved by a methylation sensitive restriction enzyme or an isoschizomer of a methylation sensitive restriction enzyme, such that the overhang on the nucleic acid of interest will base pair with the 3' or 5' single stranded end of the adaptor under appropriate conditions. The conditions will vary depending on the sequence composition (GC vs AT), the length, and the type of nucleic acid (see Sambrook et al., Molecular Cloning: a Laboratory Manual, 2nd Ed.; Cold Spring Harbor Laboratory Press, Plainview, NY, 1998).

Following the ligation of the oligonucleotide, the nucleic acid of interest is amplified using a primer complementary to the oligonucleotide. Specifically, the term "primer" as used herein refers to a sequence comprising two or more deoxyribonucleotides or ribonucleotides, preferably more than three, and more preferably more than eight, wherein the sequence is capable of initiating synthesis of a primer extension product, which is substantially complementary to a nucleic acid such as an adaptor or a ligated oligonucleotide. Environmental conditions conducive to synthesis include the presence of nucleoside triphosphates and an agent for polymerization,

such as DNA polymerase, and a suitable temperature and pH. The primer is preferably single stranded for maximum efficiency in amplification, but may be double stranded. If double stranded, the primer is first treated to separate its strands before being used to prepare extension products. In one embodiment, the primer is an oligodeoxyribo-nucleotide. The primer must be sufficiently long to prime the synthesis of extension products in the presence of the inducing agent for polymerization. The exact length of primer will depend on many factors, including temperature, buffer, and nucleotide composition. The oligonucleotide primer typically contains 12-20 or more nucleotides, although it may contain fewer nucleotides.

Primers of the invention are designed to be "substantially" complementary to each strand of the oligonucleotide to be amplified and include the appropriate G or C nucleotides as discussed above. This means that the primers must be sufficiently complementary to hybridize with their respective strands under conditions which allow the agent for polymerization to perform. In other words, the primers should have sufficient complementarity with a 5' and 3' oligonucleotide to hybridize therewith and permit amplification of CpG containing nucleic acid sequence.

Primers of the invention are employed in the amplification process which is an enzymatic chain reaction that produces exponential quantities of target locus relative to the number of reaction steps involved (e.g., polymerase chain reaction or PCR). Typically, one primer is complementary to the negative (-) strand of the locus and the other is complementary to the positive (+) strand. Annealing the primers to denatured nucleic acid followed by extension with an enzyme, such as the large fragment of DNA Polymerase I (Klenow) and nucleotides, results in newly synthesized + and - strands containing the target locus sequence. Because these newly synthesized sequences are also templates, repeated cycles of denaturing, primer annealing, and extension results in exponential production of the region (i.e., the target locus sequence) defined by the primer. The product of the chain reaction is a discrete

nucleic acid duplex with termini corresponding to the ends of the specific primers employed.

The oligonucleotide primers of the invention may be prepared using any suitable method, such as conventional phosphotriester and phosphodiester methods or automated embodiments thereof. In one such automated embodiment, diethylphosphoramidites are used as starting materials and may be synthesized as described by Beaucage, et al. (Tetrahedron Letters, 22:1859-1862, 1981). One method for synthesizing oligonucleotides on a modified solid support is described in U.S. Patent No. 4,458,066.

10

Where the CpG-containing nucleic acid sequence of interest contains two strands, it is necessary to separate the strands of the nucleic acid before it can be used as a template for the amplification process. Strand separation can be effected either as a separate step or simultaneously with the synthesis of the primer extension products. This strand separation can be accomplished using various suitable denaturing conditions, including physical, chemical, or enzymatic means, the word "denaturing" includes all such means. One physical method of separating nucleic acid strands involves heating the nucleic acid until it is denatured. Typical heat denaturation may involve temperatures ranging from about 80° to 105°C for times ranging from about 1 to 10 minutes. Strand separation may also be induced by an enzyme from the class of enzymes known as helicases or by the enzyme RecA, which has helicase activity, and in the presence of riboATP, is known to denature DNA. The reaction conditions suitable for strand separation of nucleic acids with helicases are described by Kuhn Hoffmann-Berling (CSH-Quantitative Biology, 43:63, 1978) and techniques for using RecA are reviewed in C. Radding (Ann. Rev. Genetics, 16:405-437, 1982).

20

When complementary strands of nucleic acid or acids are separated, regardless of whether the nucleic acid was originally double or single stranded, the separated strands are ready to be used as a template for the synthesis of additional nucleic acid strands. This synthesis is performed under conditions allowing hybridization of

30

primers to templates to occur. Generally synthesis occurs in a buffered aqueous solution, generally at a pH of about 7-9. Preferably, a molar excess (for genomic nucleic acid, usually about 108:1 primer:template) of the two oligonucleotide primers is added to the buffer containing the separated template strands. It is understood, however, that the amount of complementary strand may not be known if the process of the invention is used for diagnostic applications, so that the amount of primer relative to the amount of complementary strand cannot be determined with certainty. As a practical matter, however, the amount of primer added will generally be in molar excess over the amount of complementary strand (template) when the sequence to be amplified is contained in a mixture of complicated long-chain nucleic acid strands. a large molar excess is preferred to improve the efficiency of the process.

The deoxyribonucleoside triphosphates dATP, dCTP, dGTP, and dTTP are added to the synthesis mixture, either separately or together with the primers, in adequate amounts and the resulting solution is heated to about 90°-100°C from about 1 to 10 minutes, preferably from 1 to 4 minutes. After this heating period, the solution is allowed to cool to approximately room temperature, which is preferable for the primer hybridization. To the cooled mixture is added an appropriate agent for effecting the primer extension reaction (called herein "agent for polymerization"), and the reaction is allowed to occur under conditions known in the art. The agent for polymerization may also be added together with the other reagents if it is heat stable. This synthesis (or amplification) reaction may occur at room temperature up to a temperature above which the agent for polymerization no longer functions. Thus, for example, if DNA polymerase is used as the agent, the temperature is generally no greater than about 40°C. Most conveniently the reaction occurs at room temperature.

The agent for polymerization may be any compound or system which will function to accomplish the synthesis of primer extension products, including enzymes. Suitable enzymes for this purpose include, for example, E. coli DNA polymerase I, Klenow fragment of E. coli DNA polymerase I, T4 DNA polymerase, other available

DNA polymerases, polymerase muteins, reverse transcriptase, and other enzymes, including heat-stable enzymes (i.e., those enzymes which perform primer extension after being subjected to temperatures sufficiently elevated to cause denaturation such as Taq DNA polymerase, and the like). Suitable enzymes will facilitate combination
5 of the nucleotides in the proper manner to form the primer extension products which are complementary to each locus nucleic acid strand. Generally, the synthesis will be initiated at the 3' end of each primer and proceed in the 5' direction along the template strand, until synthesis terminates, producing molecules of different lengths. There may be agents for polymerization, however, which initiate synthesis at the 5' end and
10 proceed in the other direction, using the same process as described above.

Preferably, the method of amplifying is by PCR, as described herein and as is commonly used by those of ordinary skill in the art. However, alternative methods of amplification have been described and can also be employed. PCR techniques and
15 many variations of PCR are known. Basic PCR techniques are described by Saiki et al. (1988 Science 239:487-491) and by U.S. Pat. Nos. 4,683,195, 4,683,202 and 4,800,159, which are incorporated herein by reference.

The conditions generally required for PCR include temperature, salt, cation,
20 pH and related conditions needed for efficient copying of the master-cut fragment. PCR conditions include repeated cycles of heat denaturation (i.e. heating to at least about 95.degree. C.) and incubation at a temperature permitting primer: adaptor hybridization and copying of the master-cut DNA fragment by the amplification enzyme. Heat stable amplification enzymes like the pwo, *Thermus aquaticus* or
25 *Thermococcus litoralis* DNA polymerases are commercially available which eliminate the need to add enzyme after each denaturation cycle. The salt, cation, pH and related factors needed for enzymatic amplification activity are available from commercial manufacturers of amplification enzymes.

30 As provided herein an amplification enzyme is any enzyme which can be used for in vitro nucleic acid amplification, e.g. by the above-described procedures. Such

amplification enzymes include pwo, Escherichia coli DNA polymerase I, Klenow fragment of E. coli DNA polymerase I, T4 DNA polymerase, T7 DNA polymerase, Thermus aquaticus (Taq) DNA polymerase, Thermococcus litoralis DNA polymerase, SP6 RNA polymerase, T7 RNA polymerase, T3 RNA polymerase, T4 polynucleotide kinase, Avian Myeloblastosis Virus reverse transcriptase, Moloney Murine Leukemia Virus reverse transcriptase, T4 DNA ligase, E. coli DNA ligase or Q.beta. replicase. Preferred amplification enzymes are the pwo and Taq polymerases. The pwo enzyme is especially preferred because of its fidelity in replicating DNA.

Once amplified, the nucleic acid can be attached to a solid support, such as a membrane, and can be hybridized with any probe of interest, to detect any nucleic acid sequence. Several membranes are known to one of skill in the art for the adhesion of nucleic acid sequences. Specific non-limiting examples of these membranes include nitrocellulose (NITROPURE) or other membranes used in for detection of gene expression such as polyvinylchloride, diazotized paper and other commercially available membranes such as GENESCREEN, ZETAPROBE (Biorad), and NYTRAN. Methods for attaching nucleic acids to these membranes are well known to one of skill in the art. Alternatively, screening can be done in a liquid phase.

In nucleic acid hybridization reactions, the conditions used to achieve a particular level of stringency will vary, depending on the nature of the nucleic acids being hybridized. For example, the length, degree of complementarity, nucleotide sequence composition (e.g., GC v. AT content), and nucleic acid type (e.g., RNA v. DNA) of the hybridizing regions of the nucleic acids can be considered in selecting hybridization conditions. An additional consideration is whether one of the nucleic acids is immobilized, for example, on a filter.

An example of progressively higher stringency conditions is as follows: 2 x SSC/0.1% SDS at about room temperature (hybridization conditions); 0.2 x

SSC/0.1% SDS at about room temperature (low stringency conditions); 0.2 x SSC/0.1% SDS at about 42°C (moderate stringency conditions); and 0.1 x SSC at about 68°C (high stringency conditions). Washing can be carried out using only one of these conditions, e.g., high stringency conditions, or each of the conditions can be used, e.g., for 10-15 minutes each, in the order listed above, repeating any or all of the steps listed. However, as mentioned above, optimal conditions will vary, depending on the particular hybridization reaction involved, and can be determined empirically. In general, conditions of high stringency are used for the hybridization of the probe of interest.

10

The probe of interest can be detectably labeled, for example, with a radioisotope, a fluorescent compound, a bioluminescent compound, a chemiluminescent compound, a metal chelator, or an enzyme. Those of ordinary skill in the art will know of other suitable labels for binding to the probe, or will be able to ascertain such, using routine experimentation.

15

In one embodiment, representational difference analysis (RDA, see Lisitsyn et al., Science 259:946-951, 1993, herein incorporated by reference) can be performed on CpG-containing nucleic acid following MCA. MCA utilizes kinetic and subtractive enrichment to purify restriction endonuclease fragments present in one population of nucleic acid fragments but not in another. Thus, RDA enables the identification of small differences between the sequences of two nucleic acid populations. RDA uses nucleic acid from one population as a "tester" and nucleic acid from a second population as a "driver" in order to clone probes for single copy sequences present in (or absent from) one of the two populations. In one embodiment, nucleic acid from a "normal" individual or sample, not having a disorder such as a cell-proliferative disorder is used as a "driver," and nucleic acid from an "affected" individual or sample, having the disorder such as a cell proliferative disorder is used as a "tester." In one embodiment, the nucleic acid used as a "tester" is isolated from an individual having a cell proliferative disorder such as low grade astrocytoma,

20
25
30

anaplastic astrocytoma, glioblastoma, medulloblastoma, gastric cancer, colorectal cancer, colorectal adenoma, acute myelogenous leukemia, leukemia, lung cancer, renal cancer, breast cancer, prostate cancer, endometrial cancer and neuroblastoma. The nucleic acid used as a "driver" is thus normal astrocytes, normal glial cells, 5 normal brain cells, normal gastric cells, normal colorectal cells, normal leukocytes, normal lung cells, normal kidney cells, normal breast cells, normal prostate cells, normal uterine cells, and normal neurons, respectively. In an additional embodiment, the nucleic acid used as a "driver" is isolated from an individual having a cell proliferative disorder such as low grade astrocytoma, anaplastic astrocytoma, 10 glioblastoma, medulloblastoma, gastric cancer, colorectal cancer, colorectal adenoma, acute myelogenous leukemia, leukemia, lung cancer, renal cancer, breast cancer, prostate cancer, endometrial cancer and neuroblastoma. The nucleic acid used as a "tester" is thus normal astrocytes, normal glial cells, normal brain cells, normal gastric cells, normal colorectal cells, normal leukocytes, normal lung cells, normal kidney 15 cells, normal breast cells, normal prostate cells, normal uterine cells, and normal neurons, respectively. One of skill in the art will readily be able to identify the "tester" nucleic acid useful with to identify methylated nucleic acid sequences in given "driver" population.

20 KITS

The materials for use in the assay of the invention are ideally suited for the preparation of a kit. Therefore, in accordance with another embodiment of the present invention, there is provided a kit it useful for the detection of a cellular proliferative disorder in a subject having or at risk for said cellular proliferative disorder. 25 Invention kits include a carrier means compartmentalized to receive a sample in close confinement therein, one or more containers comprising a first container containing a reagent which modifies unmethylated cytosine and a second container containing primers for amplification of a CpG-containing nucleic acid, wherein the primers distinguish between modified methylated and nonmethylated nucleic acid, and 30 optionally, a third container containing a methylation sensitive restriction

endonuclease. Primers contemplated for use in accordance with the invention include those set forth in SEQ ID NOs: 1-50.

Carrier means are suited for containing one or more container means such as
5 vials, tubes, and the like, each of the container means comprising one of the separate
elements to be used in the method. In view of the description provided herein of
invention methods, those of skill in the art can readily determine the apportionment of
the necessary reagents among the container means. For example, one of the
container means can comprise a container containing an oligonucleotide for ligation to
10 nucleic acid cleaved by a methylation sensitive restriction endonuclease. One or more
container means can also be included comprising a primer complementary to the
oligonucleotide. In addition, one or more container means can also be included which
comprise a methylation sensitive restriction endonuclease. One or more container
means can also be included containing an isoschizomer of said methylation sensitive
15 restriction enzyme.

In another embodiment, the kit may comprise a carrier means containing one
or more container means comprising a solid support, wherein the solid support has a
nucleic acid sequence of CACNA1G as described herein immobilized on the solid
20 support. In one embodiment, the solid support is a membrane. Several membranes
are known to one of skill in the art for the adhesion of nucleic acid sequences.
Specific non-limiting examples of these membranes include nitrocellulose (Nitropure)
or other membranes used in for detection of gene expression such as
polyvinylchloride, diazotized paper and other commercially available membranes
25 such as GENESCREEN, ZETAPROBE (Biorad), and NYTRAN. The CACNA1G
sequences immobilized on the solid support can then be hybridized to nucleic acid
sequences produced by performing the MCA procedure, bisulfite PCR or other
methylation detection methods on the nucleic acids of a sample of interest in order to
determine if the nucleic acid sequences contained in the sample are methylated.

The term "oligonucleotide primer" refers to a sequence of two or more deoxyribo-nucleotides or ribonucleotides, preferably at least eight, which sequence is capable of initiating synthesis of a primer extension product that is substantially complementary to a target nucleic acid strand. The oligonucleotide primer typically
5 contains fifteen to twenty-two or more nucleotides, although it may contain fewer nucleotides if the primer is complementary, so as to specifically allow the amplification of the specifically desired target nucleotide sequence.

The oligonucleotide primers for use in the invention may be prepared using
10 any suitable method, such as conventional phosphotriester and phosphodiester methods or automated embodiments thereof. In one such automated embodiment, diethyl-phos-phoramidites are used as starting materials and may be synthesized as described by Beaucage et al., Tetrahedron Letter, 22: 1859-1862 (1981). One method for synthesizing oligonucleotides on a modified solid support is described in U.S.
15 Patent No. 4,458,066. The exact length of primer will depend on many factors, including temperature, buffer, and nucleotide composition. The primer must prime the synthesis of extension products in the presence of the inducing agent for amplification.

20 Primers used according to the method of the invention are complementary to each strand of mutant nucleotide sequence to be amplified. The term "complementary" means that the primers must hybridize with their respective strands under conditions which allow the agent for polymerization to function. In other words, the primers that are complementary to the flanking sequences hybridize with
25 the flanking sequences and permit amplification of the nucleotide sequence. Preferably, the 3' terminus of the primer that is extended has perfectly base paired complementarity with the complementary flanking strand.

The term "flanks CpG-rich regions" refers to those DNA sequences on
30 chromosome that are upstream (5') or downstream (3') to the DNA sequence to be amplified. The sequence to be amplified is preferably a CpG-rich region in a gene or

regulatory region associated with a gene. For example, when the nucleotide sequence to be amplified is double stranded, a first sequence that is 5' to the nucleotide sequence and a second sequence that is 5' to the nucleotide sequence on the complementary strand flank the CpG-rich DNA sequence.

5

The nucleotide sequences that flank nucleotide repeats, i.e., the nucleotide sequences to which the oligonucleotide primers hybridize, may be selected from among the following nucleotide sequences: SEQ ID NO:1-50.

10 In general, the primers used according to the method of the invention embrace oligonucleotides of sufficient length and appropriate sequence which provide specific initiation of polymerization of a significant number of nucleic acid molecules containing the target nucleic acid under the conditions of stringency for the reaction utilizing the primers. In this manner, it is possible to selectively amplify the specific
15 target nucleic acid sequence containing the nucleic acid of interest. Oligonucleotide primers used according to the invention are employed in any amplification process that produces increased quantities of target nucleic acid.

The invention includes antibodies immunoreactive with CACNA1G
20 polypeptide or functional fragments thereof. Antibody which consists essentially of pooled monoclonal antibodies with different epitopic specificities, as well as distinct monoclonal antibody preparations are provided. Monoclonal antibodies are made from antigen containing fragments of the protein by methods well known to those skilled in the art (Kohler, *et al.*, *Nature*, 256:495, 1975). The term antibody as used in
25 this invention is meant to include intact molecules as well as fragments thereof, such as Fab and F(ab')₂, Fv and SCA fragments which are capable of binding an epitopic determinant on CACNA1G.

(1) An Fab fragment consists of a monovalent antigen-binding fragment of an
30 antibody molecule, and can be produced by digestion of a whole antibody molecule

with the enzyme papain, to yield a fragment consisting of an intact light chain and a portion of a heavy chain.

(2) An Fab' fragment of an antibody molecule can be obtained by treating a whole
5 antibody molecule with pepsin, followed by reduction, to yield a molecule consisting of an intact light chain and a portion of a heavy chain. Two Fab' fragments are obtained per antibody molecule treated in this manner.

(3) An (Fab')₂ fragment of an antibody can be obtained by treating a whole antibody
10 molecule with the enzyme pepsin, without subsequent reduction. A (Fab')₂ fragment is a dimer of two Fab' fragments, held together by two disulfide bonds.

(4) An Fv fragment is defined as a genetically engineered fragment containing the
15 variable region of a light chain and the variable region of a heavy chain expressed as two chains.

(5) A single chain antibody ("SCA") is a genetically engineered single chain molecule
20 containing the variable region of a light chain and the variable region of a heavy chain, linked by a suitable, flexible polypeptide linker.

As used in this invention, the term "epitope" refers to an antigenic determinant
on an antigen, such as a CACNA1G polypeptide, to which the paratope of an
antibody, such as an CACNA1G -specific antibody, binds. Antigenic determinants
usually consist of chemically active surface groupings of molecules, such as amino
25 acids or sugar side chains, and can have specific three-dimensional structural characteristics, as well as specific charge characteristics.

As is mentioned above, antigens that can be used in producing CACNA1G
-specific antibodies include CACNA1G polypeptides or CACNA1G polypeptide
30 fragments. The polypeptide or peptide used to immunize an animal can be obtained

by standard recombinant, chemical synthetic, or purification methods. As is well known in the art, in order to increase immunogenicity, an antigen can be conjugated to a carrier protein. Commonly used carriers include keyhole limpet hemocyanin (KLH), thyroglobulin, bovine serum albumin (BSA), and tetanus toxoid. The coupled
5 peptide is then used to immunize the animal (e.g., a mouse, a rat, or a rabbit). In addition to such carriers, well known adjuvants can be administered with the antigen to facilitate induction of a strong immune response.

The invention also provides a method for detecting a cell proliferative disorder
10 associated with CACNA1G in a subject, comprising contacting a target cellular component suspected of having a CACNA1G associated disorder, with a reagent which reacts with or binds to CACNA1G and detecting CACNA1G. The target cell component can be nucleic acid, such as DNA or RNA, or it can be protein. When the component is nucleic acid, the reagent is typically a nucleic acid probe or PCR primer.
15 When the cell component is protein, the reagent is typically an antibody probe. The target cell component may be detected directly in situ or it may be isolated from other cell components by common methods known to those of skill in the art before contacting with a probe. (See for example, Maniatis, et al., Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory, N.Y. 1989; Current Protocols in
20 Molecular Biology, 1994, Ed. Ausubel, et al., Green Publ. Assoc. & Wiley Interscience.) Detection methods include Southern and Northern blot analyses, RNase protection, immunoassays and other detection assays that are known to those of skill in the art.

25 The probes can be detectably labeled, for example, with a radioisotope, a fluorescent compound, a bioluminescent compound, a chemiluminescent compound, a metal chelator, or an enzyme. Those of ordinary skill in the art will know of other suitable labels for binding to the probes or will be able to ascertain such, using routine experimentation.

Since the present invention shows that a decreased level of CACNA1G transcription is often the result of hypermethylation of the CACNA1G gene, it is often desirable to directly determine whether the CACNA1G gene is hypermethylated. In particular, the cytosine rich areas termed "CpG islands" which lie in the 5' regulatory regions of genes are normally unmethylated. The term "hypermethylation" includes any methylation of cytosine which is normally unmethylated in the CACNA1G gene sequence can be detected by restriction endonuclease treatment of CACNA1G polynucleotide (gene) and Southern blot analysis for example. Therefore, in a method of the invention, when the cellular component detected is DNA, restriction endonuclease analysis is preferable to detect hypermethylation of the CACNA1G gene. Any restriction endonuclease that includes CG as part of its recognition site and that is inhibited when the C is methylated, can be utilized. Methylation sensitive restriction endonucleases such as BssHIII, MspI, NotI or HpaII, used alone or in combination are examples of such endonucleases. Other methylation sensitive restriction endonucleases will be known to those of skill in the art. In addition, PCR can be utilized to detect the methylation status of the CACNA1G gene. Oligonucleotide primers based on any coding sequence region in the CACNA1G sequence are useful for amplifying DNA by PCR. CACNA1G is described here for exemplary purposes. The other genes described herein as being

For purposes of the invention, an antibody or nucleic acid probe specific for CACNA1G may be used to detect the presence of CACNA1G polypeptide (using antibody) or polynucleotide (using nucleic acid probe) in biological fluids or tissues. Oligonucleotide primers based on any coding sequence region in the CACNA1G sequence are useful for amplifying DNA, for example by PCR. Any specimen containing a detectable amount of CACNA1G polynucleotide or CACNA1G polypeptide antigen can be used. Nucleic acid can also be analyzed by RNA in situ methods which are known to those of skill in the art. A preferred sample of this invention is tissue of heart, renal, brain, colon, breast, urogenital, uterine, hematopoietic, prostate, thymus, lung, testis, and ovarian. Preferably the subject is human.

Various disorders which are detectable by the method of the invention include astrocytoma, anaplastic astrocytoma, glioblastoma, medulloblastoma, colon cancer, lung cancer, renal cancer, leukemia, breast cancer, prostate cancer, endometrial cancer and neuroblastoma.

5

Monoclonal antibodies used in the method of the invention are suited for use, for example, in immunoassays in which they can be utilized in liquid phase or bound to a solid phase carrier. In addition, the monoclonal antibodies in these immunoassays can be detectably labeled in various ways. Examples of types of immunoassays which
10 can utilize monoclonal antibodies of the invention are competitive and non-competitive immunoassays in either a direct or indirect format. Examples of such immunoassays are the radioimmunoassay (RIA) and the sandwich (immunometric) assay. Detection of the antigens using the monoclonal antibodies of the invention can be done utilizing immunoassays which are run in either the forward, reverse, or
15 simultaneous modes, including immunohistochemical assays on physiological samples. Those of skill in the art will know, or can readily discern, other immunoassay formats without undue experimentation.

The term "immunometric assay" or "sandwich immunoassay", includes
20 simultaneous sandwich, forward sandwich and reverse sandwich immunoassays. These terms are well understood by those skilled in the art. Those of skill will also appreciate that antibodies according to the present invention will be useful in other variations and forms of assays which are presently known or which may be developed in the future. These are intended to be included within the scope of the present
25 invention.

Monoclonal antibodies can be bound to many different carriers and used to detect the presence of CACNA1G. Examples of well-known carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and
30 modified celluloses, polyacrylamides, agaroses and magnetite. The nature of the

carrier can be either soluble or insoluble for purposes of the invention. Those skilled in the art will know of other suitable carriers for binding monoclonal antibodies, or will be able to ascertain such using routine experimentation.

5 In performing the assays it may be desirable to include certain "blockers" in the incubation medium (usually added with the labeled soluble antibody). The "blockers" are added to assure that non-specific proteins, proteases, or anti-heterophilic immunoglobulins to anti-CACNA1G immunoglobulins present in the experimental sample do not cross-link or destroy the antibodies on the solid phase
10 support, or the radiolabeled indicator antibody, to yield false positive or false negative results. The selection of "blockers" therefore may add substantially to the specificity of the assays described in the present invention.

 It has been found that a number of nonrelevant (i.e., nonspecific) antibodies of
15 the same class or subclass (isotype) as those used in the assays (e.g., IgG1, IgG2a, IgM, etc.) can be used as "blockers". The concentration of the "blockers" (normally 1-100 $\mu\text{g}/\mu\text{l}$) may be important, in order to maintain the proper sensitivity yet inhibit any unwanted interference by mutually occurring cross reactive proteins in the specimen.

20

 In using a monoclonal antibody for the in vivo detection of antigen, the detectably labeled monoclonal antibody is given in a dose which is diagnostically effective. The term "diagnostically effective" means that the amount of detectably labeled monoclonal antibody is administered in sufficient quantity to enable detection
25 of the site having the CACNA1G antigen for which the monoclonal antibodies are specific. The concentration of detectably labeled monoclonal antibody which is administered should be sufficient such that the binding to those cells having CACNA1G is detectable compared to the background. Further, it is desirable that the detectably labeled monoclonal antibody be rapidly cleared from the circulatory system
30 in order to give the best target-to-background signal ratio.

As a rule, the dosage of detectably labeled monoclonal antibody for in vivo diagnosis will vary depending on such factors as age, sex, and extent of disease of the individual. The dosage of monoclonal antibody can vary from about 0.001 mg/m.sup.2 to about 500 mg/m.sup.2, preferably 0.1 mg/m.sup.2 to about 200 mg/m.sup.2, most preferably about 0.1 mg/m.sup.2 to about 10 mg/m.sup.2. Such dosages may vary, for example, depending on whether multiple injections are given, tumor burden, and other factors known to those of skill in the art.

For in vivo diagnostic imaging, the type of detection instrument available is a major factor in selecting a given radioisotope. The radioisotope chosen must have a type of decay which is detectable for a given type of instrument. Still another important factor in selecting a radioisotope for in vivo diagnosis is that the half-life of the radioisotope be long enough so that it is still detectable at the time of maximum uptake by the target, but short enough so that deleterious radiation with respect to the host is minimized. Ideally, a radioisotope used for in vivo imaging will lack a particle emission, but produce a large number of photons in the 140-250 keV range, which may be readily detected by conventional gamma cameras.

For in vivo diagnosis, radioisotopes may be bound to immunoglobulin either directly or indirectly by using an intermediate functional group. Intermediate functional groups which often are used to bind radioisotopes which exist as metallic ions to immunoglobulins are the bifunctional chelating agents such as diethylenetriaminepentaacetic acid (DTPA) and ethylenediaminetetraacetic acid (EDTA) and similar molecules. Typical examples of metallic ions which can be bound to the monoclonal antibodies of the invention are .sup.111 In, .sup.97 Ru, .sup.67 Ga, .sup.68 Ga, .sup.72 As, .sup.89 Zr, and .sup.201 Tl.

A monoclonal antibody useful in the method of the invention can also be labeled with a paramagnetic isotope for purposes of in vivo diagnosis, as in magnetic resonance imaging (MRI) or electron spin resonance (ESR). In general, any

conventional method for visualizing diagnostic imaging can be utilized. Usually gamma and positron emitting radioisotopes are used for camera imaging and paramagnetic isotopes for MRI. Elements which are particularly useful in such techniques include ¹⁵⁷Gd, ⁵⁵Mn, ¹⁶²Dy, ⁵²Cr, and ⁵⁶Fe.

5

The present invention also provides a method for treating a subject with a cell proliferative disorder associated with of CACNA1G comprising administering to a subject with the disorder a therapeutically effective amount of reagent which modulates CACNA1G expression. In brain, breast and renal cancer cells, for example, the CACNA1G nucleotide sequence is under-expressed as compared to expression in a normal cell, therefore, it is possible to design appropriate therapeutic or diagnostic techniques directed to this sequence. Thus, where a cell-proliferative disorder is associated with the expression of CACNA1G associated with malignancy, nucleic acid sequences that modulate CACNA1G expression at the transcriptional or translational level can be used. In cases when a cell proliferative disorder or abnormal cell phenotype is associated with the under expression of CACNA1G, for example, nucleic acid sequences encoding CACNA1G (sense) could be administered to the subject with the disorder.

The term "cell-proliferative disorder" denotes malignant as well as non-malignant cell populations which often appear to differ from the surrounding tissue both morphologically and genotypically. Such disorders may be associated, for example, with absence of expression of CACNA1G. Essentially, any disorder which is etiologically linked to expression of CACNA1G could be considered susceptible to treatment with a reagent of the invention which modulates CACNA1G expression.

The term "modulate" envisions the suppression of methylation of CACNA1G polynucleotide when CACNA1G is under-expressed. When a cell proliferative disorder is associated with CACNA1G expression, such methylation suppressive reagents as 5-azacytadine can be introduced to a cell. Alternatively, when a cell

30

- proliferative disorder is associated with under-expression of CACNA1G polypeptide, a sense polynucleotide sequence (the DNA coding strand) encoding CACNA1G polypeptide, or 5' regulatory nucleotide sequences (i.e., promoter) of CACNA1G in operable linkage with CACNA1G polynucleotide can be introduced into the cell.
- 5 Demethylases known in the art could also be used to remove methylation.

- The present invention also provides gene therapy for the treatment of cell proliferative disorders which are mediated by CACNA1G. Such therapy would achieve its therapeutic effect by introduction of the appropriate CACNA1G
- 10 polynucleotide which contains a CACNA1G structural gene (sense), into cells of subjects having the proliferative disorder. Delivery of sense CACNA1G polynucleotide constructs can be achieved using a recombinant expression vector such as a chimeric virus or a colloidal dispersion system.

- 15 The polynucleotide sequences used in the method of the invention may be the native, unmethylated sequence or, alternatively, may be a sequence in which a nonmethylatable analog is substituted within the sequence. Preferably, the analog is a nonmethylatable analog of cytidine, such as 5-azacytadine. Other analogs will be known to those of skill in the art. Alternatively, such nonmethylatable analogs could
- 20 be administered to a subject as drug therapy, alone or simultaneously with a sense structural gene for CACNA1G or sense promoter for CACNA1G operably linked to CACNA1G structural gene.

- In another embodiment, a CACNA1G structural gene is operably linked to a
- 25 tissue specific heterologous promoter and used for gene therapy. For example, a CACNA1G gene can be ligated to prostate specific antigen (PSA)-prostate specific promoter for expression of CACNA1G in prostate tissue. Other tissue specific promoters will be known to those of skill in the art. Alternatively, the promoter for a tumor suppressor gene can be linked to the CACNA1G structural gene and used for
- 30 gene therapy.

Various viral vectors which can be utilized for gene therapy as taught herein include adenovirus, herpes virus, vaccinia, or, preferably, an RNA virus such as a retrovirus. Preferably, the retroviral vector is a derivative of a murine or avian retrovirus. Examples of retroviral vectors in which a single foreign gene can be inserted include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), and Rous Sarcoma Virus (RSV). Most preferably, a non-human primate retroviral vector is employed, such as the gibbon ape leukemia virus (GaLV), thereby providing a broader host range than murine vectors, for example.

10

A number of additional retroviral vectors can incorporate multiple genes. All of these vectors can transfer or incorporate a gene for a selectable marker so that transduced cells can be identified and generated. Retroviral vectors can be made target specific by inserting, for example, a polynucleotide encoding a sugar, a glycolipid, or a protein. Preferred targeting is accomplished by using an antibody to target the retroviral vector. Those of skill in the art will know of, or can readily ascertain without undue experimentation, specific polynucleotide sequences which can be inserted into the retroviral genome to allow target specific delivery of the retroviral vector containing the CACNA1G sense or antisense polynucleotide.

20

Since recombinant retroviruses are defective, they require assistance in order to produce infectious vector particles. This assistance can be provided, for example, by using helper cell lines that contain plasmids encoding all of the structural genes of the retrovirus under the control of regulatory sequences within the LTR. These plasmids are missing a nucleotide sequence which enables the packaging mechanism to recognize an RNA transcript for encapsidation. Helper cell lines which have deletions of the packaging signal include but are not limited to .PSI.2, PA317 and PA12, for example. These cell lines produce empty virions, since no genome is packaged. If a retroviral vector is introduced into such cells in which the packaging

25

signal is intact, but the structural genes are replaced by other genes of interest, the vector can be packaged and vector virion produced.

Another targeted delivery system for CACNA1G polynucleotide is a colloidal dispersion system. Colloidal dispersion systems include macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. The preferred colloidal system of this invention is a liposome. Liposomes are artificial membrane vesicles which are useful as delivery vehicles in vitro and in vivo. It has been shown that large unilamellar vesicles (LUV), which range in size from 0.2-4.0 μm can encapsulate a substantial percentage of an aqueous buffer containing large macromolecules. RNA, DNA and intact virions can be encapsulated within the aqueous interior and be delivered to cells in a biologically active form (Fraley, et al., Trends Biochem. Sci., 6:77, 1981). In addition to mammalian cells, liposomes have been used for delivery of polynucleotides in plant, yeast and bacterial cells. In order for a liposome to be an efficient gene transfer vehicle, the following characteristics should be present: (1) encapsulation of the genes of interest at high efficiency while not compromising their biological activity; (2) preferential and substantial binding to a target cell in comparison to non-target cells; (3) delivery of the aqueous contents of the vesicle to the target cell cytoplasm at high efficiency; and (4) accurate and effective expression of genetic information (Mannino, et al., Biotechniques, 6:682, 1988).

The composition of the liposome is usually a combination of phospholipids, particularly high-phase-transition-temperature phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on pH, ionic strength, and the presence of divalent cations.

Examples of lipids useful in liposome production include phosphatidyl compounds, such as phosphatidylglycerol, phosphatidylcholine, phosphatidylserine,

phosphatidylethanolamine, sphingolipids, cerebroside, and gangliosides. Particularly useful are diacylphosphatidylglycerols, where the lipid moiety contains from 14-18 carbon atoms, particularly from 16-18 carbon atoms, and is saturated. Illustrative phospholipids include egg phosphatidylcholine, dipalmitoylphosphatidylcholine and distearoylphosphatidylcholine.

The targeting of liposomes has been classified based on anatomical and mechanistic factors. Anatomical classification is based on the level of selectivity, for example, organ-specific, cell-specific, and organelle-specific. Mechanistic targeting can be distinguished based upon whether it is passive or active. Passive targeting utilizes the natural tendency of liposomes to distribute to cells of the reticulo-endothelial system (RES) in organs which contain sinusoidal capillaries. Active targeting, on the other hand, involves alteration of the liposome by coupling the liposome to a specific ligand such as a monoclonal antibody, sugar, glycolipid, or protein, or by changing the composition or size of the liposome in order to achieve targeting to organs and cell types other than the naturally occurring sites of localization.

The surface of the targeted delivery system may be modified in a variety of ways. In the case of a liposomal targeted delivery system, lipid groups can be incorporated into the lipid bilayer of the liposome in order to maintain the targeting ligand in stable association with the liposomal bilayer. Various linking groups can be used for joining the lipid chains to the targeting ligand.

In general, the compounds bound to the surface of the targeted delivery system will be ligands and receptors which will allow the targeted delivery system to find and "home in" on the desired cells. A ligand may be any compound of interest which will bind to another compound, such as a receptor.

In general, surface membrane proteins which bind to specific effector molecules are referred to as receptors. In the present invention, antibodies are preferred receptors. Antibodies can be used to target liposomes to specific cell-surface ligands. For example, certain antigens expressed specifically on tumor cells, referred to as tumor-associated antigens (TAAs), may be exploited for the purpose of targeting CACNA1G antibody-containing liposomes directly to the malignant tumor. Since the CACNA1G gene product may be indiscriminate with respect to cell type in its action, a targeted delivery system offers a significant improvement over randomly injecting non-specific liposomes. Preferably, the target tissue is human brain, colon, breast, lung, and renal origin. A number of procedures can be used to covalently attach either polyclonal or monoclonal antibodies to a liposome bilayer. Antibody-targeted liposomes can include monoclonal or polyclonal antibodies or fragments thereof such as Fab, or F(ab').sub.2, as long as they bind efficiently to an antigenic epitope on the target cells. Liposomes may also be targeted to cells expressing receptors for hormones or other serum factors.

It should be noted that as used herein and in the appended claims, the singular forms "a," "and," and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a cell" includes a plurality of such cells and reference to "the restriction enzyme" includes reference to one or more restriction enzymes and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs. Although any methods, devices and materials similar or equivalent to those described herein can be used in the practice or testing of the invention, the preferred methods, devices and materials are now described.

All publications mentioned herein are incorporated herein by reference in full for the purpose of describing and disclosing the methodologies which are described in the publications which might be used in connection with the presently described

invention. The publications discussed above and throughout the text are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the inventors are not entitled to antedate such disclosure by virtue of prior invention.

5

The following example is intended to illustrate but not to limit the invention in any manner, shape, or form, either explicitly or implicitly. While they are typical of those that might be used, other procedures, methodologies, or techniques known to those skilled in the art may alternatively be used.

10

EXAMPLE 1
IDENTIFICATION OF CACNA1G AS A TARGET FOR
HYPERMETHYLATION ON HUMAN CHROMOSOME 17q21

15 In order to isolate genes differentially methylated in cancer cells as opposed to non-cancerous cells the following experimental protocols were used. An example of the results obtained is provided hereinabove in the description of the isolation and characterization of human CACNA1G.

20 **Tissue Samples and Cell Lines**

Forty-nine primary colorectal cancers, 28 colorectal adenomas, 16 primary gastric cancers and 17 acute myelogenous leukemia samples were used for methylation analyses. DNA from eight colon cancer cell lines (Caco2, RKO, SW48, HCT116, DLD1, Lovo, SW837, HT29), 4 lung cancer cell lines (OH3, H249, H157, H209), 4 glioblastoma cell lines (Dauy, D283, U87, U373), 8 breast cancer cell lines (MB-468, MCF7, MB-231, MB-474, MB-435, MB-453, BT20, CAMA1, SKBR3), 7 hematopoietic tumor cell lines (CEM, Raji, KG1A, HL60, ML-1, MoIt3, K562), and 4 prostate cancer cell lines (DU145, DUPRO, LNCAP, TSUPRL) were also investigated. DNA was extracted by standard procedures. RNA was isolated from cell lines and adenomas using TRIZOL (GIBCO-BRL). For re-expression analysis,

30

cell lines were treated with 5-Aza-deoxycytidine (SIGMA) at a final concentration of 1 M for 6 days. All tissue samples were obtained from patients who gave informed consent according to institutional guidelines.

5 RT-PCR.

Six g of total RNA, was reverse transcribed using the SUPERScript kit (GIBCO-BRL) for first strand cDNA synthesis. One hundred ng of cDNA was used as template for RT-PCR reactions. To design the RT-PCR primers, Blast search was performed using the rat Cacna1G cDNA sequence (Genbank AF027984) reported previously (25) and exon-intron boundaries of the human CACNA1G were predicted by this analysis. Each primer set was designed to amplify the cDNA across several introns. Primer sequences and PCR conditions are available, at <http://www.med.jhu.edu/methylation/primers>. GAPDH was also amplified as a control using primers GAPDHF: 5'-CGGAGTCAACGGATTGGTCGTAT-3' (SEQ ID NO:53) and GAPDHR: 5'-AGCCTTCTCCATGGTGGTGAAGAC-3' (SEQ ID NO:54). All reactions were performed with RT(-) controls. PCR amplification was performed for 35 cycles of 95° C 30 sec, 60-65°C for 30 sec, 72°C for 30 sec, and the products were analyzed by agarose gel electrophoresis.

20 DNA Sequencing and Data Analysis.

PCR reaction products were precipitated with ethanol, resuspended in diluted water and cloned into the pCR2.1 vector using the TA cloning kit (Invitrogen) according to the manufacturer's instruction. After transformation, plasmid DNA was purified using the Wizard Miniprep Kit (Promega). DNA sequence analysis was carried out at the Johns Hopkins University Sequence Facility using automated DNA sequencers (Applied Biosystems). Sequence homology was identified by the BLAST program of the National Center for Biological Information (NCBI) available at <http://www.ncbi.nlm.nih.gov/BLAST/>. An IMAGE cDNA clone (Genbank: H13333) was identified by BLAST analysis using the sequence of BAC AC004590 (Genbank) which includes MINT31. Putative genes (G1 and G2) were identified by GENSCAN

(available at <http://ccr-081.mit.edu/GENSCANMIT.html>) using the BAC sequence data. IMAGE cDNA clone H1333) was then obtained from the American Type Culture Collection and completely sequenced. Potential transcription factor binding sites and promoter prediction were examined using the TESS and TSSG programs
5 respectively, available at the Baylor College of Medicine BCM Launcher (http://kiwi.imgen.bcm.tmc.edu:8088/search_launcher/launcher.html/). The nucleotide sequence of part of the 5' end of the cDNA of CACNA1G has been submitted to Genbank.

10 Bisulfite-PCR Methylation Analysis.

Bisulfite treatment was performed as reports previously. (Herman, J. G., Graff, J. R., Myohanen, S., Nelkin, B. D. and Baylin, S.B. *Proc. Natl. Acad. Sci. USA*. 93:9821-9826, 1996). Briefly, 2 g of genomic DNA was denatured with 2 M NaOH for 10 min., followed by incubation with 3 M Na-bisulfite, pH 5.0 for 16 hours at
15 50°C. After treatment DNA was purified using a Wizard Miniprep Column (Promega), precipitated with ethanol and resuspended in 20 l of diluted water. Two l of the aliquot was used as template for each PCR reaction. Semi-quantitative bisulfite-PCR was performed essentially as described. Xiong, Z. and Laird, P. W. COBRA: a sensitive and quantitative DNA methylation assay. *Nucleic Acids Res.*
20 25:2532-2534, 1997. To avoid overestimation of the methylated alleles, the following points were considered. First, primers were designed to contain a minimum number of CpG dinucleotides in the sequence to avoid the biased amplification of methylated alleles. If primers do contain CpG sites, they were designed to amplify methylated and unmethylated alleles equally (using a mixture of C or T for sense and a mixture of
25 G or A for antisense primers). Second, the primers were designed to contain a maximum number of thymidines converted from cytosines to avoid amplifying the non-converted genomic sequence. Third, restriction sites which only appear after bisulfite conversion (e.g. ACGC to ACGT) were used (regions 1-8). PCR was performed as described previously (Herman, supra). Primer sequences, annealing
30 temperature and PCR cycles are available at <http://www.jhu.edu/methylation/primers>.

Twenty % of the PCR products were digested with the appropriate restriction enzymes, precipitated with ethanol and separated by 5% polysacrylamide gel electrophoresis. Gels were stained with ethidium bromide, and the intensity of each allele was calculated by densitometry, using the Image Quant software (Molecular
5 Dynamics).

Although the invention has been described with reference to the presently preferred embodiment, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention is
10 limited only by the following claims.

What is claimed is:

1. An isolated nucleic acid molecule comprising the coding region for a T-type calcium channel and regulatory sequences associated therewith.
2. The nucleic acid molecule of claim 1, wherein said associated regulatory sequences contain CpG-rich regions.
3. The nucleic acid molecule of claim 2, wherein the state of methylation of the CpG-rich regions is determinative of the presence of a cellular proliferative disorder in a subject from which the nucleic acid molecule is isolated.
4. The nucleic acid molecule of claim 2, wherein hypermethylation of said CpG islands is indicative of the presence of a cellular proliferative disorder in a subject from which said nucleic acid is isolated.
5. The nucleic acid molecule of claim 1, wherein said T-type calcium channel is CACNA1G.
6. The nucleic acid molecule of claim 5, wherein said nucleic acid comprises the nucleic acid sequence of SEQ ID NO: 51.
7. The nucleic acid molecule of claim 6, wherein one or more of regions 1-8 comprises methylated bases.
8. A substantially purified polypeptide encoded by the polynucleotide of SEQ ID NO:51.
9. The polypeptide of claim 8, wherein the polypeptide has an amino acid sequence as set forth in SEQ ID NO:52.

10. A method for detecting a cellular proliferative disorder in a subject comprising:
 - a) contacting a nucleic acid-containing specimen from the subject with an agent that provides a determination of the methylation state of at least one gene or associated regulatory region of the gene; and
 - b) identifying aberrant methylation of regions of the gene or regulatory region, wherein aberrant methylation is identified as being different when compared to the same regions of the gene or associated regulatory region in a subject not having said cellular proliferative, thereby detecting a cellular proliferative disorder in the subject.
11. The method of claim 10, wherein the regions of said gene are contained within CpG rich regions.
12. The method of claim 10, wherein the gene is selected from the group consisting of APOB, CACNA1G, CDX2, EGFR, FBN1, GPR37, HSPA6, IQGAP2, KL, PAR2, PITX2, PTCH, SDC1, SDC4 and combinations thereof.
13. The method of claim 10, wherein aberrant methylation comprises hypermethylation when compared to the same regions of the gene or associated regulatory regions in a subject not having the cellular proliferative disorder.
14. The method of claim 13, wherein the regions comprise regulatory regions of CACNA1G.
15. The method of claim 14, wherein the regions comprise regions 1-8 of CACNA1G.
16. The method of claim 15, wherein the regions comprise regions 1-2 of CACNA1G.
17. The method of claim 15, wherein the regions comprise regions 5-7 of CACNA1G.

18. The method of claim 15, wherein the regions comprise regions 4 and 8 of CACNA1G.
19. The method of claim 10, wherein the agent is a pair of primers that hybridize with a target sequence in the gene or associated regulatory region of the gene.
20. The method of claim 19, wherein the primers hybridize with a target polynucleotide sequence having the sequence selected from the group consisting of SEQ ID NO:55-103 and SEQ ID NO:104.
21. The method of claim 20, wherein the primers are in consecutive pairs selected from the group consisting of SEQ ID NO:1-49 and SEQ ID NO:50.
22. The method of claim 10, wherein the nucleic acid-containing specimen comprises a tissue selected from the group consisting of brain, colon, urogenital, lung, renal, prostate, pancreas, liver, esophagus, stomach, hematopoietic, breast, thymus, testis, ovarian, and uterine.
23. The method of claim 10, wherein the nucleic acid-containing specimen is selected from the group consisting of serum, urine, saliva, blood, cerebrospinal fluid, pleural fluid, ascites fluid, sputum, stool, and biopsy sample.
24. The method of claim 10, wherein said cellular proliferative disorder is selected from the group consisting of low grade astrocytoma, anaplastic astrocytoma, glioblastoma, medulloblastoma, gastric cancer, colorectal cancer, colorectal adenoma, acute myelogenous leukemia, lung cancer, renal cancer, leukemia, breast cancer, prostate cancer, endometrial cancer and neuroblastoma.

25. A kit useful for the detection of a cellular proliferative disorder in a subject comprising:
- a) carrier means compartmentalized to receive a sample therein;
 - b) one or more containers comprising a first container containing a reagent which modifies unmethylated cytosine and a second container containing primers for amplification of a CpG-containing nucleic acid, wherein the primer hybridizes with a target polynucleotide sequence having the sequence selected from the group consisting of SEQ ID NO:55-103 and SEQ ID NO:104.
26. The kit of claim 25, further comprising a third container containing a methylation sensitive restriction endonuclease.
27. The kit of claim 25, wherein said modifying reagent is bisulfite.
28. The kit of claim 25, wherein the primers are selected from the group consisting of SEQ ID NO:1-49 and SEQ ID NO:50.
29. Isolated oligonucleotide primer(s) for detection of a methylated CpG-containing nucleic acid wherein the primer hybridizes with a target polynucleotide sequence having the sequence selected from the group consisting of SEQ ID NO:55-103 and SEQ ID NO:104.
30. The primers of claim 29, wherein said primers are selected from the group consisting of SEQ ID NO:1-49 and SEQ ID NO:50.
31. An isolated nucleic acid molecule having at least one methylated Cytosine of a CpG dinucleotide in a CpG-rich region and encoding a gene selected from the group consisting of APOB, CACNA1G, CDX2, EGFR, FBN1, GPR37, HSPA6, IQGAP2, KL, PAR2, PITX2, PTCH, SDC1, and SDC4.

32. The nucleic acid molecule of claim 31, wherein the methylated C residue of a CpG dinucleotide is located within a CpG-rich region selected from the group consisting of SEQ ID NO:105-118 and SEQ ID NO:119.

Figure 1A




Figure 1B




Figure 2

Methylation pattern	Island 1				R3 M	R4		Island 2				R8 T B	Expression of <i>hCACNA1G</i>
	R1		R2			M T		R5		R6	R7		
	M	B	T	B		M	T	B	T	H	E		
1													++
2													+++
3													+++
4													0
5													0-++

FIGURE 3A (SEQ ID NO:51) CACNA1G nucleotide sequence

CCTTTTCGTTTCGCCCTCTCGGGGCGGCTTCGCCGAAGGTAGCGCCGAATCCGGCAACCGGAGCCTGGGCGCGAAGCGAAG
AAGCCGGAACAAAGTGAGGGGGAGCCGGCCGGCTGGCCCGGAAGCCCCAGGGGCGCAGGGGAAGCGGGACTCGCGCCGG
GCGGGGTTTCCCTGCGCCCCGCGCGCCCCGCGGGCAGCATGCCCCCTGCGGGCAGGGGGAGCTGGGCTGAACTGGCCCTCCC
GGGGGCTCAGCTTGCGCCCTAGAGCCCAACAGATGTGCCCCCGCGGGGCCCCCGGGTTGCGTGAGGACACCTCCTCTGA
GGGGCGCGCTTGCCCTCTCCGATCGCCCGGGGCCCGGCTGGCCAGAGGATGGACGAGGAGGAGATGGAGCGGGCG
CCGAGGAGTCGGGACAGCCCCGGAGCTTCATGCGGCTCAACGACCTGTGCGGGGCGGGGGCCGGCCGGGGCCGGGGTCA
GCAGAAAAGGACCCGGGCGCGGACTCCGAGGCGGAGGGGTGCCGTACCCGGCGCTGGCCCCGGTGGTTTTTCTTCTA
CTTGAGCCAGGACAGCCGCCCGCGGAGCTGGTGCTCCGACAGGTCTGTAAACCCCTGGTTTGAGCGCATCAGCATGTTGG
TCATCCTTCTCAACTGCGTGAACCTGGGCATGTTCCGGCCATGCGAGGACATCGCCTGTGACTCCAGCGCTGCCGGATC
CTGCAGGCCTTTTGATGACTTCATCTTTGCCCTTTTGCCGATGGAGATGGTGGTGAAGATGGTGGCTTGGGCATCTTTGG
GAAAAAGTGTTTACCTGGGAGACACTTGAACCGGCTTGACTTTTTTCATCGTCATCGCAGGGATGCTGGAGTACTCGCTGG
ACCTGCAGAACGTCAGCTTCTCAGCTGTGAGACAGTCCGTGTGCTGCGACCGCTCAGGGCCATTAAACGGGTGCCAGC
ATGCGCATCCTTTGTACGTTGCTGTGATACGCTGCCATGCTGGGCAACGTCCTGCTGCTGCTTCTTCTGCTTCTTCTT
CATCTTCGGCATCGTCGGCSTCCAGCTGTGGGCGAGGGTGTCTTCGGAACCGATGCTTCTACCTGAGAATTTACAGCCTCC
CCCTGAGCGTGACCTGGAGCGCTATTACCAGACAGAGAACGAGGATGAGAGCCCTTCATCTGCTCCCAGCCACGCGAG
AACGGCATCGGTCCTGCAGAACGCTGCCCCACGCTGCGCGGGGACGGGGCGGTGGCCACCTTGCGGTCTGGACTATGA
GGCTACAACAGCTCCAGCAACACCACCTGTGTCAACTGGAACCACTACTACCACTGCTCAGCGGGGAGCAACAAC
CCTTCAAGGGCGCCATCAACTTTGACAACATTGGCTATGCTGATCGCCATCTTCCAGGTATCAGCGTGGAGGGCTGG
GTCGACATCATGTACTTTGTGATGGATGCTCATTCTTCTACAATTTTCTACTTCTATCCTCCTCATCATCGTGGGCTC
CTTCTTCATGATCAACCTGTGCTGCTGGTGTATGCCACGAGTCTCAGAGACCAAGCAGCGGAAAGCCAGCTGATGC
GGGAGCAGCGTGTGCGGTTCTGTCCAACGCCAGCACCTTGGCTAGCTTCTCTGAGCCCGGAGCTGCTATGAGGAGCTG
CTCAAGTACCTGGTGATACCTTCGTAAGGACGCCCGCAGGCTGGCTCAGGTCTCTCGGGCAGCAGGTGTGCGGGTTGG
GCTGCTCAGCAGCCAGCACCCCTCGGGGCGCAGGAGACCCAGCCAGCAGCTGCTCTCGCTCCCACCGCCGCTAT
CCGCTCCACCACTGGTGCAACCAACCAACCATCACCACCACTACCACCTGGGCAATGGGACGCTCAGGGCCCCCGG
GCCAGCCCGGAGATCCAGGACAGGGATGCCAATGGGTCCCGCGGCTCATGTGCCACCACTCGACGCTGCCCTCTC
CGGGGCCCCCTTGGTGGCGCAGAGTCTGTGCACAGCTTCTACCATGCCAGTGCCTTAGAGCCAGTCCGCTGCCAGG
CGCCCCCTCCCAGGTCCCCATCTGAGGCATCCGGCAGGACTGTGGGCGAGCGGAAGGTGTATCCACCGTGACACACAGC
CCTCCACCGGAGACGCTGAAGGAGAAGGCACTAGTAGAGGTGGCTGCCAGCTCTGGGCCCCCAACCTCACCAGCCTCAA
CATCCCACCGGGCCCTACAGCTCCATGCACAAGCTGCTGGAGACACAGGTACAGGTGCCTGCCAAAGCTCTTGCAAGA
TCTCCAGCCCTTGCTTGAAAGCAGACAGTGGAGCCTGTGCTCCAGACAGCTGCCCTACTGTGCCGGGCGGGGCGAGG
GAGGTGGAGCTCGCCGACCGTGAATGCCTGACTCAGACAGCGAGGAGTTTATGAGTTACACAGGATGCCACAGCACAG
CGACCTCCGGGACCCCCACAGCCGGCGGCAACGGAGCTGGGCCAGATGCAGAGCCAGCTCTGTGCTGGCCTTCTGGA
GGCTAATCTGTGACACCTTCCGAAAGATTGTGGACAGCAAGTACTTTGGCCGGGAATCATGATCGCCATCCTGGTCAAC
ACACTCAGCATGGGCATCGAATACCAGAGCAGCCCGAGGAGCTTACCAACGCCCTAGAAATCAGCAACATCGTCTTCAC
CAGCCTCTTTGCCCTGGAGATGCTGCTGAAGCTGCTTGTGTATGGTCCCTTTGGCTACATCAAGAATCCCTACAACATCT
TCGATGGTGTCAATTGTGGTTCATCAGCGTGTGGGAGATCGTGGGCCAGCAGGGGGCGGCCTGTGGTGTGCGGACCTTC
CGCTGATGCGTGTGCTGAAGCTGGTGGCTTCTGCGGCGCTGCAGCGGAGCTGGTGGTGTCTATGAAGACCATGGA
CAACGTGGCCACCTTCTGCATGCTGCTTATGCTCTTCTCATCTTCTAGCATCTTGGGCATGCATCTCTTCGGCTGCA
AGTTTGCCTCTGAGCGGATGGGGACACCTGCCAGACCGGAAGAATTTGACTCCTTGCTCTGGGCCATCGTCACTGTC
TTTCAGATCCTGACCAGGAGGACTGGAACAAAGTCTCTACAATGGTATGGCTCCACGTGCTCCTGGGCGGCCCTTTA
TTTCATTGCCCTCATGACCTTCGGCACTACGTGCTCTTCAATTTGCTGGTGGCCATTCTGGTGGAGGGCTTCCAGGCGG
AGGGAGATGCCAACAAAGTCCGAATCAGAGCCCCGATTTCTTCTACCCAGCCTGGATGGTGTATGGGGACAGGAAGAAGTGC
TTGGCCTTGGTGTCCCTGGGAGAGCACCCGAGCTGCGGAAGAGCCTGCTGCCGCTCTCATCATCCACACGGCCGCCAC
ACCCATGTGCTGCCAAGAGCACCAAGCAGCGGCTGGGGAGGCGCTGGGCCCTGCGTGCAGCCGACCAAGCAGCAGCG
GGTGGCAGAGCCTGGGGCGGCCACGAGATGAAGTACCGCCAGCGCCCGCAGCTCTCCGACAGCCCTGGAGCGCT
GCAAGCAGCTGGACCAGCAGGCGCTCCAGCCGGAACAGCCTCGGCCGTGACCCAGCCTGAAGCGGAGAAGCCAAAGTGG
AGAGCGGCGGTCCCTGTTGTGGGAGAAGGCCAGGAGAGCAGGATGAAGAGGAGAGCTCAGAAGAGGAGCGGGCCAGCC
CTGCGGGCAGTGACCATCGCCACAGGGGGTCCCTGGAGCGGGAGGCCAAGAGTTCTTTGACCTGCCAGACACACTGCAG
GTGCCAGGGCTGCATCGCACTGCCAGTGGCCGAGGGTCTGCTTCTGAGCACCAGGACTGCAATGGCAAGTGGCTTCAGG
GCGCCTGGCCCCGGCCCTGCGGCTGATGACCCCCACTGGATGGGGATGACGCCGATGACGAGGGCAACCTG

Figure 3B (SEQ ID NO:52)

MDEEDGAGAEESGQPRSFMRNLNDLSGAGGRPGPGSAEKDPGSA
DSEAEGLPYPALAPVVFYLSQDSRPRSWCLRTVCNPWFERISMLVILLNCVTLGMR
PCEDIACDSQRCRILQAFDDFIFAFFAVEMVVKMVALGIFGKKCYLGDTWNRLDFFIV
IAGMLEYSLDLQNVSPSAVRTVRVLRPLRAINRVPSMRILVTLLLDTLPLGNVLLLC
FFVFFIFGIVGVQLWAGLLRNRCFLPENFSLPLSVDLERYQTENEDESFFICSQPRE
NGMRSCRSVPTLRGDGGGPPCGLDYEAYNSSNTTCVNWNQYYTNC SAGEHNPFKGA
INFDNIGYAWIAIFQVITLEGWVDIMYFVMDAHSFYNFIFYFILLIIVGSFFMINLCLV
VIATQFSETKQRESQLMREQRVRFSLNASTLASFPSEPGSCYEELLKYLVIYILRKAARR
LAQVSRAAGVRVGLLSSPAPLGGQETQPSSSCSRSHRRLSVHHLVHHHHHHHHHHYHLG
NGTLRAPRASPEIQDRDANGSRRLMLPPPSTPALSGAPPGGAESVHSFYHADCHLEPV
RCQAPPRSPSEASGRTVGSGKVYPTVHTSPPPETLKEKALVEVAASSGPPTLTSLNI
PPGPYSSMHKLLLETQSTGACQSSCKISSPCLKADSGACGPDSCPYCARAGAGEVELAD
REMPDSDSEAVYEFTQDAQHSDLRDPHSRRQRS LGPDAEPSSVLAFWRLICDTFRKIV
DSKYFGRGIMIAILVNTLSMGIYHEQPEELTNAL EISNIVFTSLFALEMLLKL LVYG
PFGYIKNPYNIFDGVIVVISVWEIVGQGGGLSVLRTFRLMRVLKLVRF PALQRQLV
VLMKTMNDNVATFCMLLM LFI FIFSI LGMHLFGCKFASERDGD TLPDRKNFDSLLWAI V
TVFQILTQEDWNKVLYNGMASTSSWAALYFIALMTFGNYVLFNLLVAILVEGFQAEGD
ANKSESEPDDFFSPSLDGDGDRKKCLALVSLGEHPELRKSLLPPLIHTAATPMSLPKS
TSTGLGEALGPASRR TSSSGSAEPGAAHEMKSPPSARSSPHSPWSAASSWTSRRSSRN
SLGRAPSLKRRSPSGERRSLLSGEGQESQDEEESSEEERAS PAGSDHRHRGSLEREAK
SSFDL PDTLQVPGLHRTASGRGSASEHQDCNGKSASGRLARALRPDDPPLDGDDADDE
GNL

FIGURE 4A

APOB CpG ISLAND (SEQ ID NO:105)

ccccggaggcgccctttggaccttttgaatcctggcgctcttcagcctgggcttctataaatggggtcgggcgccggccgcgcattc
ccaccgggagctcgggggtgagtgcccttctcgggttgctgcccgtgaggagccccagccagccaggcgccgagggccgaggcc
aggccgcagcccaggagccgcccaccgcagctggcgatggaccgccgaggcccgctgctggcgctgcctgcgtgctgctgct
gctgctggcgggcgccaggccgggtgagtgccgcccgtctgcccgcagcagaggagcgggaggagccggcgaccgaggtt
ggccggggcagcctgggctaggccagaggaggcagccacagggtccaggggcagtggggggattggaccagctggcgcccc
tgaggctcaggatggggggcgccggatggaggggctgaggagggggtctccggagcctgcctccctcctgaaagggtgaaacctgtgc
cgggtggtccccctgtcgggcccctagcaccgcgtgggaagacgtgggaagctc

CACNA1G CpG ISLAND (SEQ ID NO:106)

cctcgggccctacgccaggaccccgccgaatactctgattcttcgggctccctccaaggagtgcccaaagaccccaatggccaatagg
aaagtgggttcgggtcggcagcagctgattggctccagccttcgggagcggaccaggggcaaggggaggggagaggggcggtcct
gggttttggggtgggaatcggattccagctgtggttctccttcgctcccggcactgccacggcgagccaatggcgcgagg
ctcggggccggcggtccggcgattggctgcggggctgtcggggcgggggccgaggctgaagtgaagtgaggatccagctgtg
gtgtgcgcggggctcctgcgcggcgttgcctgcctcgcctcgcctcgcggcgaggaggaggtgtggcgccggcgacagctacg
gcagcgccagccaccgcggcggtgcggcgggcgatccgcctccactcccgggactgccccactgtctccccgcccctc
ccggacagtgagcccgcgcgggggcgaggagccgccccacccctccaagcccacccctaaagagatccctctccctcc
cccgccgctggcgaggagccgggacgatgctgaccccttagatccggctccagctgcgcggcgagggaaggggcgccccctcccg
gacccccgcccctccgctgcccccttttctgctccctcctggggcggttcgccgaaggtagcggcgaatccggcaaccggagcct
gggcgcgaaagcgaagaagccggaacaaagtgagggggagccggcggtggccgggaagccccaggggcgaggggaagcgg
gactgcgcggggcggggtttccctgcgccccggcgcccccgggcgagcatcccctgcgggcagggggagctgggctgaactggc
cctcccgggggtcagcttgcgccttagagcccaccagatgtcccccgccggggccccgggttgcgtgaggacacctctctgagg
Ggcgcgcgttgcctctccggatcggcgggggccccgggtggccagaggatggacgaggaggaggtgagcggcgccgagga
gtcgggacagccccggagctcatgcggctcaacgacctgtcggggggccggggggcgccggggcggggtcagcagaaaaggaccc
gggcagcgcggaactccgagcgagggggtgcccgtaccggcgctggccccgggtggttttcttacttagccaggacagccccc
cggagctggTgtctccgcacgggtgttaaccatactctcggggcacgacggccagggcgcggggtcagaagggggacggcgccac
cgccgggggtcgggggggaagaagaccaccgcccaggtgagtcgaagttagcccgagggttaggcggatggggggggggctgcc
agggaggggaggggggcaccagagtgggagcggagacgcgagcaggtctcgtcgttaaccggggttaccacactgcgtacacaca
cctcagctctcctgggttgggggggtgggatccaggccagagaagagagctgtgccccgctggctcgcagctggacgcccctcagat
gtgtcaggggagggtcgtatcctccagatgtgggaagcttcgggagcctgggagctgtactctgccccgcggggttagcgagctgggt
ttggttccgagtttgggggggtgggtggggcggtggggagggaagctcgggggacggaggaggggggaccgcaatctcctgggtt
tccctcttccccgccccaaagtgttcggcggttctagatgttggggggcggggaccaggtcctggccacctcaccctccctgcg
gggtggaggcacaacaaggagattccggcgggcggtgatgtcagggcgcgagaatgagaacaagatgtggtggaggggagctgtctgc
ccccggagctgggagtgaggcccccttccgctagagcccagtgccgcgggtgcctcctaccgatctccattcgatgc

FIGURE 4B

CDX2 CpG Island (SEQ ID NO:107)

ctcgtaatcacggaagccggcctggggctccgcacgccagcctgtgcgggtcttccccgcctctgcagcctagtgggaaggaggt
gggaggaagaaggaagaaaggaggaggaggaggcaggccagaggaggaccgcctcggaggcagaagagccgcgagg
agccagcggagcaccgcgggctggggcgcagccaccgccgctcctcaggtccctcggcccttcccttcgtgccccggcagcctc
cagcgtcgtccccaggcagcatggtgaggtctgtcccggtccctcggccaccatgtacgtgagcta

EGFR CpG Island (SEQ ID NO:108)

gtccgcggggaccgggtccagaggggcagtgctgggaacgcccctctcgaaattaactcctcagggcaccgctccccccatgcgcc
gccccactcccgcggagactaggtcccgcggggccaccgtgtccaccgcctcggggccgctggccttgggtccccgctgtgtgttct
cctccctcctcctcgtatctcctcctcctctgtcctcccgatccctcctcggccgctgtccctcctcctccccccctgcctccgcgcctc
ggcccgcgagctagacgtccgggcagccccggcgagcgccggccgcagcagcctcctcccccgacgggtgtgagcggccggc
gcggcgaggcgccggagctccgagctagccccgcggcgccggccggccagaccggacgacaggccacctcgtcgcgtccggccg
agtccccgcctcggcgcaacgccacaaccaccgcgcagggccccctgactccgtccagtattgatcgggagagccggagcgagctctt
cggggagcagcgtacgacctccgggacggcgggggcagcgctcctggcgctgctggctgcgtctgccCggcgagtcgggctctg
Gaggaaaagaaggttaaggcggtgtctcgcggctccccgcggccccggatcgcggcccgacccccgagccccgccaaccgcacc
gcgcaccggcttcggcgccccgccccgcttctcgttcttctgagatcacgtgcggccggaccgggaccggggagggaacggg
acgtttcgttctcggcggggagagctctggggcgggcgaggaggagacgcgtgggacaccgggctgcaggccaggcggggaacgg
ccggcgggacctccggcgccccgaaccgctcccaacttctcctcacttccccgcccagctgcgcaggatggcgctcagtgggcgaa
agccgggtgctggtggcgccctggggcggggtcccgacgggctccccgcgctgttctccaggcgcgacggggtcctggcgcg
acccgagggcgctgccaccggcgagactgcctgttaggggaagctgagggaaggaacccaaaaatacagcctccgctcggacccg
cgggacaggcggttctgagaggacctccccgcctccgctccgcgaggtctcaactgaagccggcgcccgccagcctggcccc
ggccctctccaggctccccgcgatcctcgttccccagtggtgagtcgcagcctcgacctgggagctgggagaactcgtctaccaccactg
cggctcccggggagggtggtgctggcggttagttcctcgttggcaaaaggcaggtggggtccgaccgccccttggcgcgagac
ccggccgctcgcctcggcggtgcgcctcgttgcctatccaagagtccccccactcccgggacccagctcctccgcgcccgc
gccgaaagccccaggctctcctcgtatggcgccctcgcggagacgtccgggtctgtccacctgcagcccttcggtcgcgttgggctt
gcgggtggagcgggacgggctgtccggccactgcaggggggagtcgggactcttgagcggaagccccg

FIGURE 4C

FBN1 CpG Island (SEQ ID NO:109)

agagccgcgtctggagtgggctctcgacacccagggaagtgggggcggcagagccctctctcggcggcacagcagccctctccgcgggtccgg
cctgcgacgcgccagcttagcctcccgccctccggcgtctgctgagtgccggcgggagaggcgagggagcgctaccgggagggcggggc
agcggggactggtttctctcggccaggccctccggggcaaccgtctccagcgcgacattcttggtgcaggtggaacagcttctgctccggtagggtt
cacctatcgcgggagaggtaatctcggatctaaacctcgagcccgagagcgggctaaaaccgctactccacctcttccatttctccctccacctc
aagacaaaaagtcaggccggcgaggacctgatcacctctgctcctccactcgctaatctcgagcgagaggccccgaccgagggcgagg
ctgcaaaaggagtggaaggaggatggatggggccgggggtgggtgggtgatgagggcgacgaaggagggggtgtcatttcttttcttttt
ttaaaaaagtatttctctcgagaaaccgtcgcgggacgatacttgaagaggtgggaaaggagggggtgctggggagcccgcgagagactgtg
gggtccacaagcgagacaggagccacagctgGgacagctcgagcgaggagccgagcagtggttagcgccacgactgggagcagccgcccg
ctctctcgggagtcggagcccgcttctccagtgggtgcagccgggtccgacgggggtcggcgccaccggggctggagctcgggccacgga
ggcttttgcgttgcggcnngaggcgaggacaggagctgggtgaggggtgtcccggaacgtccaaactggnctgagaccttccctgcct
gacagcttctgncggggtcttgggtccggnccggcgtagatgtcggggggcgggtgcacgccccggagtcggcggggacggcgcggtgctt
ccagctggcgagaggcgagctgaggagtggggcggttcagagcgcgcatcgcgcaattcgtccgctaaaaataaaaccagagagctcgc
ccggggcttaggaccgctggggatatgggtactttgcgcgcgcttcttggcgggggccgggagggcgagggttggccggggctgtcgccggg
gcctgggctttccagccagctgtggaccaaacggtcttcccttaccaaattaactgcgcacagggcgccgacnggttgggctttgggaatgggacc
gcgagcttcagcatccgatgccctgaaagtctccccgcctcggggaattgtctgtgttcagcttggcagggcgccgctgaagtgggagcagcgct
ggagaaggcgggagggagcccgccgggggacggcgcgggatagcgggaccccgcgcgcggtgcgcttcaggcgagcgggcgccg
cagaccgagccccggcgcggaaggcgcgggagccggtggcgctcggcatcatgctcagggcgctgtgtggagatcgccctgggatt
accgtgcttttagctgtctacacgagccatggggcgggacgccaatttgagggtgggaactgaaggaaaccagagccagtcgggccaagagaaga
ggcgggtggagacacgacgcgttaaagggtaaaggaaaccggttccctc

GPR37 (SEQ ID NO:110)

Tccgccccgcacccgccctagccccgggctcggggacctgtcagggtggtttcgacagctggggaattaacctgtcccgccatccctagcctcag
ccgcgagggctccgcgctccgcttgttccctccagctctccgagtggaagcgctacaaatggcttgaatgaacgtgtgtgggttagtgagtg
gtgaaccaccaggggatccgctctcccaaaaccagtatctctccgaggaggagggcgaaggagtgggaggaggcaacgagccgagagtcgagctt
cgcgggcgcgcgagcggtgagcgggggcgaggcgggccacctcccttcccgccgcgactgcctggcccgcgcggttccaggca
ccaccttcccgtccgggtgagcccgctgtggcagtgactgctcccgcggtagcggcactgtccaccgacgagcgggcgccctcttcccccttct
ccccacgatttcttctcgcggcgacgacctccagcagcctgcttccccgctcactttagctggaggagaagcaacttggcagtgcccg
ggggttgaatcccgttctcctcgagcagtaggctcgcaagtcgctgggttaggtggggcaagagtttccggcgcatcagcgctgcttcggac
tggttgcaacgtgttccagcgagctgggagcggggtgtgactgcgagtcgtctggggagggggacttgttttcttctctagacacctcggcttga
actgatcaaacgctgtcgaaa

HSPA6 CpG Island (SEQ ID NO:111)

tgatttcgatggtaacatatcttgggtcttccgctgggctctcagcgccctccaaggcagcccgagggcggtgctcgcctcagggatcctccac
agccccggggagaccttgctctaaagtgtgttgcagctctgcccacaaccgcgctcctcagagccagccgggagggagctagaaccttccccgc
gtttcttcagcagccctgagtcagaggcggtggtgcttgaagtacggcccgccagccttcttgggtctacggaccgatccgcccgaaccttccccg
ggtcagcgccgcgctgcgcggcggtgactcagccggcgggcgggcgaggctctcagctggcggggaagggtcgggaaagggttcgcg
cgcggggtcggggaggtgcaaaaggatgaaaagcccggtggagcgagctgagcagatccggcggggtggcgggcagagaaaccgcaggagag
cctcactgtgagcggccctcgacggggcgagcagcctcgtggcctcagcatccgacaa

FIGURE 4D

IQGAP2 CpG Island (SEQ ID NO:112)

Agagttcacttttacttcagtgtcagcgcgcggcgccgtggctggctctggcgagagagcaccgagggagtggtcgcagatcttcgg
gcggctaggggaaatcggcgagagggcgccgatccgagcgcgccggcgggcgagagcccgagcctggccagcgagggtagcc
gcggggggcgccccggcgggccccggagacgcgcaggatgccacacgaagagctgccgtcgtcagagaccccgctatgg
ctctattgtggacgatgaaa

KL CpG Island (SEQ ID NO:113)

ctcgaaagagggcgcggtggcgcgctctccccgcgagcatctcacctaagggggaatcccttcagcgcacggcggaagtccccctc
ggctgtcccacctggcagtcctctaggtttcgccagtccttaattggctccagcaatgtccagccggagcttcttggccctccgagtgg
gagaaaagttagagcaggtgcttccccagcgcgctccgctagggcccgagcatcccgccccaagtgggaaaagttggctg
gcgccttttctccccagcgaagccgctccagggctgctctcagaggacgcgcggcaggcaagagaatgaacctgagcgtccacgaaac
gtcctgcacggctcccgaggctgggagaaacaggtgcttctccgacgtccgcggcgacgcctgccgcaccttggccgctgccgcg
ccccccccggcacccctcggcgccccctgccccacccccagtgccagggcgaggcagtcggctcgcaggtattattgc
cagcggagccccggggagcgggggtggcgcgccggcggtggcgGgcggcgcgcgggggcgcgggcataaagggcgcg
ggcgcgggggccccggagcctggctcccgcgagcatgccgccagcgccccggcgcccgccggccgcccgcgctgctgt
cgctgctgctggtgctgctgggcctggcgggccgcccgtgcgtgcggagccggcgacggcgcgagacctgggcccgtttctcg
gcctcctcccccgaggccgcgggcctctccagggcaccttccccgacggcttctctgggccgtgggcagcgccgctaccagaccg
agggcggtggtggcagcagcacggcaaggggtgctccatctgggacacgttaccaccacccccctggcacccccgggagactcccgga
acgccagtctgctgtggcgccccgctccgctgcagcccgccaccggggacgtagccagcgacagctacaacaacgttctccgga
cacggaggcgctgcgcgagctcgggtcactcactaccgttctccatctcgtggcgcgagtgctcccaatggcagcgcgggcgctcc
ccaaccgcgaggggctgcgtactaccggcgctgctggagcggtgcggagctggcggtgcagccgtggtaccctgtaccctg
ggacctgccccagcgctgcagacgctacggcggtggccaaccgcgccctggccgaccacttcagggtattacgggagctctgc
ttccgccacttcggcggtcaggtcaagtactggatcaccatcgacaacccctacgtggtggcctggcacggctacgccaccggcgccctg
gccccggcatccggggcagcccgcggtcggtacctggtggcgcaaa

PAR2 CpG Island (SEQ ID NO:114)

Cccggggcggtggcctcccgaggtgagtagctgctccttcggtttccctgaaacctaacccgccctggggagggcgcgagcagaggt
ccgattcggggcaggtgagaggctgactttctcgggtcgtccagtgagctctgagttcgaatcgcgggcgggattccccgcgcg
ccggcgtcggggcttcaggaggatcgggagccccagcgcggtggctgctgggggcccgcctctgc

PITX2 CpG Island (SEQ ID NO:115)

PTCA CpG Island (SEQ ID NO:116)

[illegible]

CGCGCCGCGGCACTGTCTGCCCGTGCCCCCTGCCCTGAACCTTCTTCTCCTGCGCCCCCTGCCCCCTAT
TTGCAGCCTAAACTCCTGTACGGCTGCCACATTTCTTAACATCTTGGAGGGGAGGCGGAGTGAGAG
AGGCGGAGAGAGGAAGGGGGGAGGGAGCCGAAATAAAGGTGGTTTCTTTTTTGGCAGCCAGTTTTG
GTTTTGTTGAGCATGAAATCTCTGCTCCCTTAAAAAATTATTCTCGGAAAAAGATATCCCCCCCCGTTTT
CCAGGTTTTGAGCCGCCTCTCCTTAGGGCCTGGTCGGGGGAGGAAAAGTTGTAAACAAATTGCCACCT
TAAATTGCGGGTGCGANTCTGCGGAGCTGCCGGGTTTATTGTGTTACGAGGCTCGCTGAAATGTGTG
GAATCCAGGGAAGGCGAGCACCCAGACGGGGGGCCCGGGGGCCGCGGCCAGCGCCGGGGAAATGC
CGCGCCGGGGAGCAGCATGCGCCGGCCTGAGCCCTTCCCTTTGCACTCGGCTGTTTTTTACGTTTAACC
AGAAAGGAAGGGAGAGGAGGGAAGATCCATGTGGCTGCCCTCTCCGATCACAAATATTGTCGTA
GTTGCACTGGCTGCCCANTTCTAATTCAGCTCACACAGCNTNTCCCCACGCTATGGAAATGCGTCG
GGAGTGAACTCCGGCGGCCGC

FIGURE 4F

SDC1 CpG Island (SEQ ID NO:118)

Ggagaggtgcgggccgaatccgagccgagcagaggaatccggcagtagagagcggactccagccggcggaccctgcagccctcg
cctgggacagcggcgcgctgggcaggcgcccaagagagcatcgagcagcgggaaccgcgaagccggcccgagccgcgacccgc
gcagcctgccgctctcccgccgggtccgggcagcatgaggcgcgcgcgctctggctctggctgtgcgcgctggcgctgagcctgc
agctggccctgccgca

SDC4 CpG Island (SEQ ID NO:119)

Agtaggagccggcgggctcgggcaggcggggtcccttggggttccaactccggcgggcgggcgagtgccccgcaggcctcgcttc
actggggaattccggcggggtgcggcgggcgggcgggcgggcgggcgggcgggcgggcgggcgggcgggcgggcgggcgggcg
cgccccccggggccactcgccgcagcctgcgcgccttctccagtcgcggtgccatggccccgcccgtctgttcgcgctgctgtt
cttcgtaggcggagtcgccgagtcgggtgggtgcttggaggttccgggctggggcggaagcggggcgagggccggtgcctccttgtt
cgtcggagcgtgggatggggggggcagatcgggggtacgtaccccaaccggacaccgaggccccgggaaactttgttgaaacttt
gctccggggtcacgggcccagcctccgggatggcttcacgcgcgtgcggcctcgctgttcttcccgctccccgggcctcagccc
cgccgcgggctacgggctcgttagtgactaagccggtgtcaactcttcaactccacaccctcgtcccttccctggtagccctggggcagg
cttgagcgctgaatccctctcgtctcggggcgcccagagcagacagctttaggatccgagatggccctgggggtcggggggctgc
gtgtactcggaagggggagggttttagggtgtgcgaggccc

FIGURE 5

MINT31 (SEQ ID NO:120)

CCCGGGGCCT	CTATCCTGGC	GGGAAGGGCA	GGCCGACCCG	GCAGACTGCG	GCCTCTCGGG
AGGGAAGAAG	GTGTCAGACG	CGCGGAGCAA	CCATAAATAG	CCCCCCTTTC	CCAGAAGACG
GCACGGGGTT	CAAGACTCAG	GCGCCGCATA	CTCAGAATGA	GAGCAGAGAC	TCCCCGCCAGG
AAAAAAGGGC	ACTTAGGGGA	TCTGCTCATT	AACATGAAAT	GCAAATGAGC	CCGCCCCGGCC
TCATTTACAC	AACTCTGTGC	ATGGATTCCG	CGAAAGGGCA	ACCAGGGAGA	CGACGGCGCA
GCAGCCACTC	TGCCACTTCC	CCCATCCCCT	CCCCCCCATC	GGCCGGGGCG	GGAAGTGA
CGACCCCAAC	CCTCTGCGGC	GGCGGGAGGT	GCGCGGGGGC	TGCGTGGGTG	GTGCAGCCTT
AGGGGAGTGA	ACAACGCCCA	GGGGTGATGG	CCTCAGCAAA	GTGAGGGGTG	GTGATGGAGG
TCATCCGACC	CATCCCGCCG	CCTCTCCGCA	GTGGCGCAAG	CGCCCCAAAA	TCTCCGGAGA
NGGAACTGAG	TGACCCACTA	GGTTCCGCCG	TGTCTACCTC	TCGCAGATGT	TGGGGAAGTG
CTTCCCGGCG	TCTAATCCTC	GCTGTTCCCC	CCTCCACCGG	CGCCCAGCAC	ACCCGCGGCG
CTCCGCTCCC	GGG				

SEQUENCE LISTING

<110> The Johns Hopkins University School of Medicine

<120> CACNA1G POLYNUCLEOTIDE POLYPEPTIDE AND
METHODS OF USE THEREFOR

<130> JHU1590WO

<140> 09/398,522

<141> 1999-09-15

<160> 120

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Bisulfite-PCR primer

<400> 1

gtttttgtag tttgggtttt tt

22

<210> 2

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Bisulfite-PCR primer

<221> misc_feature

<222> (0)...(0)

<223> r = G or A

<400> 2

rcacaacaacr ccaacaac

18

<210> 3

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Bisulfite-PCR primer

<221> misc_feature

<222> (0)...(0)

<223> y = C or T

<400> 3

ttgtggygtt ggygatagtt

20

<210> 4

<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 4
acraaaaaaaaaa tctctt

26

<210> 5
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 5
gggggygttt tttttyggat ttt

23

<210> 6
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 6
ttccctacr cccctaaaac ttcc

24

<210> 7
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 7
gtaggttaga gggagggaty gtt

23

<210> 8

<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<400> 8
aaaacaaacc tcaccatact acct 24

<210> 9
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 9
tttgatttyg tttagtattg at 22

<210> 10
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<400> 10
cccttacctt tcttttcct 19

<210> 11
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 11
tttttttatt gygtaattt tg 22

<210> 12
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<400> 12

tttccccacc tcttcaaata 20

<210> 13
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<400> 13
ggttaggtgg ggtaagagtt t 21

<210> 14
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 14
aacrtttaat ccaattacaa acc 23

<210> 15
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<400> 15
tttttttagta gttttgagtt agagg 25

<210> 16
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<400> 16
tttcatcctt ttacacctcc c 21

<210> 17
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<400> 17
tttattttta ttttagtggt ag 22

<210> 18
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 18
ctcttcttat aacatcctac
20

<210> 19
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 19
gtagtatggt ygtagygtt t
21

<210> 20
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<400> 20
aaatacccta aaaaaaaccc
20

<210> 21
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 21
yggttttttt gaaatttaat tc
22

<210> 22
<211> 19
<212> DNA

<213> Artificial Sequence

<220>

<223> Bisulfite-PCR primer

<221> misc_feature

<222> (0)...(0)

<223> r = G or A

<400> 22

aactccrcaat cctcctaaa

19

<210> 23

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Bisulfite-PCR primer

<221> misc_feature

<222> (0)...(0)

<223> y = C or T

<400> 23

taagtgttty gttaggtttt tt

22

<210> 24

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Bisulfite-PCR primer

<400> 24

ccaaactcca ctacacaata ac

22

<210> 25

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Bisulfite-PCR primer

<221> misc_feature

<222> (0)...(0)

<223> y = C or T

<400> 25

gttggtttgt taatyggagt

20

<210> 26

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 26
ttaccaaccr aaaccatatt

20

<210> 27
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<400> 27
aatgtgtgga atttaggga

19

<210> 28
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<400> 28
taaaacaacc aactacaact tac

23

<210> 29
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 29
agaggaatty ggtagtagag ag

22

<210> 30
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<400> 30
cacaaccaa accaaaac

18

<210> 31
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<400> 31
ggggatttgt ttggtagtgg 20

<210> 32
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Bisulfite-PCR primer

<400> 32
cccgaaattc cccaataaa 19

<210> 33
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 33
gayggygtag tagttatttt gtt 23

<210> 34
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<400> 34
catcaccacc cctcacttta c 21

<210> 35
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 35

tttgggtatt tatagttttt tggag

25

<210> 36

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<221> misc_feature

<222> (0)...(0)

<223> r = G or A

<400> 36

aatctaccrc cttcactcac tc

22

<210> 37

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<221> misc_feature

<222> (0)...(0)

<223> y = C or T

<400> 37

tttaggagyg ttaatgtgag gtt

23

<210> 38

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<400> 38

ctaaaaaac ccaatcttaa aaaaac

26

<210> 39

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<400> 39

tggataaagg atgtttgggg tttg

24

<210> 40

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<400> 40

ccctcccctt acccctaaat cc

22

<210> 41

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<221> misc_feature

<222> (0)...(0)

<223> y = C or T

<400> 41

aatyggattt tagttgtggt tttt

24

<210> 42

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<400> 42

cacaccacaa ctaaaccct cact

24

<210> 43

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<221> misc_feature

<222> (0)...(0)

<223> y = C or T

<400> 43

ttgtggygtt ggygatagtt

20

<210> 44

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<221> misc_feature

<222> (0)...(0)

<223> r = G or A

<400> 44

acraaaaaaaaa aaaaaaaaaa tctctt

26

<210> 45

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<221> misc_feature

<222> (0)...(0)

<223> y = C or T

<400> 45

gggggygttt tttttyggat ttt

23

<210> 46

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<221> misc_feature

<222> (0)...(0)

<223> r = G or A

<400> 46

ttcccctacr cccctaaaac ttcc

24

<210> 47

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<400> 47

gggagtttgg gagttgtatt ttggt

25

<210> 48

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<400> 48
aaccaaatta aaaaatcaaa ccctaa

26

<210> 49
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 49
gaggggggat ygtaattttt tg

22

<210> 50
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer useful for bisulfite/PCR analysis of
CACNA1G

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 50
ccraaatctc cttattatac ctccaa

26

<210> 51
<211> 3993
<212> DNA
<213> Artificial Sequence

<220>
<223> CAGNA1G - a gene encoding a T-type calcium channel

<221> CDS
<222> (373)...(3993)

<400> 51
ccttttcgtt cgccctctcg gggcggttc gccgaaggta gcgccgaatc cggcaaccgg 60
agcctgggag cgaagcgaag aagccggaac aaagtgaggg ggagccggcc ggctggcccc 120
ggaagcccca gggcgagcagg ggaagcggga ctgcgcggcg gcgggggttc cctgcgcccc 180
ggcgccccgc gggcagcatg cccctgcggg cagggggagc tgggctgaac tggccctccc 240
gggggctcag cttgcgcctt agagcccacc agatgtgccc ccgcccgggc ccccggttg 300
cgtgaggaca cctcctctga gggcgccgc ttgccctctt ccggatcgcc cggggccccg 360
gctggccaga gg atg gac gag gag gag gat gga gcg ggc gcc gag gag tcg 411
Met Asp Glu Glu Glu Asp Gly Ala Gly Ala Glu Glu Ser
1 5 10

13

gga cag ccc cgg agc ttc atg cgg ctc aac gac ctg tcg ggg gcc ggg	459
Gly Gln Pro Arg Ser Phe Met Arg Leu Asn Asp Leu Ser Gly Ala Gly	
15 20 25	
ggc cgg ccg ggg ccg ggg tca gca gaa aag gac ccg ggc agc gcg gac	507
Gly Arg Pro Gly Pro Gly Ser Ala Glu Lys Asp Pro Gly Ser Ala Asp	
30 35 40 45	
tcc gag gcg gag ggg ctg ccg tac ccg gcg ctg gcc ccg gtg gtt ttc	555
Ser Glu Ala Glu Gly Leu Pro Tyr Pro Ala Leu Ala Pro Val Val Phe	
50 55 60	
ttc tac ttg agc cag gac agc cgc ccg cgg agc tgg tgt ctc cgc acg	603
Phe Tyr Leu Ser Gln Asp Ser Arg Pro Arg Ser Trp Cys Leu Arg Thr	
65 70 75	
gtc tgt aac ccc tgg ttt gag cgc atc agc atg ttg gtc atc ctt ctc	651
Val Cys Asn Pro Trp Phe Glu Arg Ile Ser Met Leu Val Ile Leu Leu	
80 85 90	
aac tgc gtg acc ctg ggc atg ttc cgg cca tgc gag gac atc gcc tgt	699
Asn Cys Val Thr Leu Gly Met Phe Arg Pro Cys Glu Asp Ile Ala Cys	
95 100 105	
gac tcc cag cgc tgc cgg atc ctg cag gcc ttt gat gac ttc atc ttt	747
Asp Ser Gln Arg Cys Arg Ile Leu Gln Ala Phe Asp Asp Phe Ile Phe	
110 115 120 125	
gcc ttc ttt gcc gtg gag atg gtg gtg aag atg gtg gcc ttg ggc atc	795
Ala Phe Phe Ala Val Glu Met Val Val Lys Met Val Ala Leu Gly Ile	
130 135 140	
ttt ggg aaa aag tgt tac ctg gga gac act tgg aac cgg ctt gac ttt	843
Phe Gly Lys Lys Cys Tyr Leu Gly Asp Thr Trp Asn Arg Leu Asp Phe	
145 150 155	
ttc atc gtc atc gca ggg atg ctg gag tac tcg ctg gac ctg cag aac	891
Phe Ile Val Ile Ala Gly Met Leu Glu Tyr Ser Leu Asp Leu Gln Asn	
160 165 170	
gtc agc ttc tca gct gtc agg aca gtc cgt gtg ctg cga ccg ctc agg	939
Val Ser Phe Ser Ala Val Arg Thr Val Arg Val Leu Arg Pro Leu Arg	
175 180 185	
gcc att aac cgg gtg ccc agc atg cgc atc ctt gtc acg ttg ctg ctg	987
Ala Ile Asn Arg Val Pro Ser Met Arg Ile Leu Val Thr Leu Leu Leu	
190 195 200 205	
gat acg ctg ccc atg ctg ggc aac gtc ctg ctg ctc tgc ttc ttc gtc	1035
Asp Thr Leu Pro Met Leu Gly Asn Val Leu Leu Leu Cys Phe Phe Val	
210 215 220	
ttc ttc atc ttc ggc atc gtc ggc gtc cag ctg tgg gca ggg ctg ctt	1083
Phe Phe Ile Phe Gly Ile Val Gly Val Gln Leu Trp Ala Gly Leu Leu	
225 230 235	
cgg aac cga tgc ttc cta cct gag aat ttc agc ctc ccc ctg agc gtg	1131
Arg Asn Arg Cys Phe Leu Pro Glu Asn Phe Ser Leu Pro Leu Ser Val	
240 245 250	

gac ctg gag cgc tat tac cag aca gag aac gag gat gag agc ccc ttc	1179
Asp Leu Glu Arg Tyr Tyr Gln Thr Glu Asn Glu Asp Glu Ser Pro Phe	
255 260 265	
atc tgc tcc cag cca cgc gag aac ggc atg cgg tcc tgc aga agc gtg	1227
Ile Cys Ser Gln Pro Arg Glu Asn Gly Met Arg Ser Cys Arg Ser Val	
270 275 280 285	
ccc acg ctg cgc ggg gac ggg ggc ggt ggc cca cct tgc ggt ctg gac	1275
Pro Thr Leu Arg Gly Asp Gly Gly Gly Gly Pro Pro Cys Gly Leu Asp	
290 295 300	
tat gag gcc tac aac agc tcc agc aac acc acc tgt gtc aac tgg aac	1323
Tyr Glu Ala Tyr Asn Ser Ser Ser Asn Thr Thr Cys Val Asn Trp Asn	
305 310 315	
cag tac tac acc aac tgc tca gcg ggg gag cac aac ccc ttc aag ggc	1371
Gln Tyr Tyr Thr Asn Cys Ser Ala Gly Glu His Asn Pro Phe Lys Gly	
320 325 330	
gcc atc aac ttt gac aac att ggc tat gcc tgg atc gcc atc ttc cag	1419
Ala Ile Asn Phe Asp Asn Ile Gly Tyr Ala Trp Ile Ala Ile Phe Gln	
335 340 345	
gtc atc acg ctg gag ggc tgg gtc gac atc atg tac ttt gtg atg gat	1467
Val Ile Thr Leu Glu Gly Trp Val Asp Ile Met Tyr Phe Val Met Asp	
350 355 360 365	
gct cat tcc ttc tac aat ttc atc tac ttc atc ctc ctc atc atc gtg	1515
Ala His Ser Phe Tyr Asn Phe Ile Tyr Phe Ile Leu Leu Ile Ile Val	
370 375 380	
ggc tcc ttc ttc atg atc aac ctg tgc ctg gtg gtg att gcc acg cag	1563
Gly Ser Phe Phe Met Ile Asn Leu Cys Leu Val Val Ile Ala Thr Gln	
385 390 395	
ttc tca gag acc aag cag cgg gaa agc cag ctg atg cgg gag cag cgt	1611
Phe Ser Glu Thr Lys Gln Arg Glu Ser Gln Leu Met Arg Glu Gln Arg	
400 405 410	
gtg cgg ttc ctg tcc aac gcc agc acc ctg gct agc ttc tct gag ccc	1659
Val Arg Phe Leu Ser Asn Ala Ser Thr Leu Ala Ser Phe Ser Glu Pro	
415 420 425	
ggc agc tgc tat gag gag ctg ctc aag tac ctg gtg tac atc ctt cgt	1707
Gly Ser Cys Tyr Glu Glu Leu Leu Lys Tyr Leu Val Tyr Ile Leu Arg	
430 435 440 445	
aag gca gcc cgc agg ctg gct cag gtc tct cgg gca gca ggt gtg cgg	1755
Lys Ala Ala Arg Arg Leu Ala Gln Val Ser Arg Ala Ala Gly Val Arg	
450 455 460	
gtt ggg ctg ctc agc agc cca gca ccc ctc ggg ggc cag gag acc cag	1803
Val Gly Leu Leu Ser Ser Pro Ala Pro Leu Gly Gly Gln Glu Thr Gln	
465 470 475	
ccc agc agc agc tgc tct cgc tcc cac cgc cgc cta tcc gtc cac cac	1851
Pro Ser Ser Ser Cys Ser Arg Ser His Arg Arg Leu Ser Val His His	
480 485 490	

ctg gtg cac cac cac cac cac cat cac cac cac tac cac ctg ggc aat Leu Val His His His His His His His His His His Tyr His Leu Gly Asn 495 500 505	1899
ggg acg ctc agg gcc ccc cgg gcc agc ccg gag atc cag gac agg gat Gly Thr Leu Arg Ala Pro Arg Ala Ser Pro Glu Ile Gln Asp Arg Asp 510 515 520 525	1947
gcc aat ggg tcc cgc cgg ctc atg ctg cca cca ccc tcg acg cct gcc Ala Asn Gly Ser Arg Arg Leu Met Leu Pro Pro Pro Ser Thr Pro Ala 530 535 540	1995
ctc tcc ggg gcc ccc cct ggt ggc gca gag tct gtg cac agc ttc tac Leu Ser Gly Ala Pro Pro Gly Gly Ala Glu Ser Val His Ser Phe Tyr 545 550 555	2043
cat gcc gac tgc cac tta gag cca gtc cgc tgc cag gcg ccc cct ccc His Ala Asp Cys His Leu Glu Pro Val Arg Cys Gln Ala Pro Pro Pro 560 565 570	2091
agg tcc cca tct gag gca tcc ggc agg act gtg ggc agc ggg aag gtg Arg Ser Pro Ser Glu Ala Ser Gly Arg Thr Val Gly Ser Gly Lys Val 575 580 585	2139
tat ccc acc gtg cac acc agc cct cca ccg gag acg ctg aag gag aag Tyr Pro Thr Val His Thr Ser Pro Pro Pro Glu Thr Leu Lys Glu Lys 590 595 600 605	2187
gca cta gta gag gtg gct gcc agc tct ggg ccc cca acc ctc acc agc Ala Leu Val Glu Val Ala Ala Ser Ser Gly Pro Pro Thr Leu Thr Ser 610 615 620	2235
ctc aac atc cca ccc ggg ccc tac agc tcc atg cac aag ctg ctg gag Leu Asn Ile Pro Pro Gly Pro Tyr Ser Ser Met His Lys Leu Leu Glu 625 630 635	2283
aca cag agt aca ggt gcc tgc caa agc tct tgc aag atc tcc agc cct Thr Gln Ser Thr Gly Ala Cys Gln Ser Ser Cys Lys Ile Ser Ser Pro 640 645 650	2331
tgc ttg aaa gca gac agt gga gcc tgt ggt cca gac agc tgc ccc tac Cys Leu Lys Ala Asp Ser Gly Ala Cys Gly Pro Asp Ser Cys Pro Tyr 655 660 665	2379
tgt gcc cgg gcc ggg gca ggg gag gtg gag ctc gcc gac cgt gaa atg Cys Ala Arg Ala Gly Ala Gly Glu Val Glu Leu Ala Asp Arg Glu Met 670 675 680 685	2427
cct gac tca gac agc gag gca gtt tat gag ttc aca cag gat gcc cag Pro Asp Ser Asp Ser Glu Ala Val Tyr Glu Phe Thr Gln Asp Ala Gln 690 695 700	2475
cac agc gac ctc cgg gac ccc cac agc cgg cgg caa cgg agc ctg ggc His Ser Asp Leu Arg Asp Pro His Ser Arg Arg Gln Arg Ser Leu Gly 705 710 715	2523
cca gat gca gag ccc agc tct gtg ctg gcc ttc tgg agg cta atc tgt Pro Asp Ala Glu Pro Ser Ser Val Leu Ala Phe Trp Arg Leu Ile Cys 720 725 730	2571

gac acc ttc cga aag att gtg gac agc aag tac ttt ggc cgg gga atc Asp Thr Phe Arg Lys Ile Val Asp Ser Lys Tyr Phe Gly Arg Gly Ile 735 740 745	2619
atg atc gcc atc ctg gtc aac aca ctc agc atg ggc atc gaa tac cac Met Ile Ala Ile Leu Val Asn Thr Leu Ser Met Gly Ile Glu Tyr His 750 755 760 765	2667
gag cag ccc gag gag ctt acc aac gcc cta gaa atc agc aac atc gtc Glu Gln Pro Glu Glu Leu Thr Asn Ala Leu Glu Ile Ser Asn Ile Val 770 775 780	2715
ttc acc agc ctc ttt gcc ctg gag atg ctg ctg aag ctg ctt gtg tat Phe Thr Ser Leu Phe Ala Leu Glu Met Leu Leu Lys Leu Leu Val Tyr 785 790 795	2763
ggc ccc ttt ggc tac atc aag aat ccc tac aac atc ttc gat ggt gtc Gly Pro Phe Gly Tyr Ile Lys Asn Pro Tyr Asn Ile Phe Asp Gly Val 800 805 810	2811
att gtg gtc atc agc gtg tgg gag atc gtg ggc cag cag ggg ggc ggc Ile Val Val Ile Ser Val Trp Glu Ile Val Gly Gln Gln Gly Gly Gly 815 820 825	2859
ctg tcg gtg ctg cgg acc ttc cgc ctg atg cgt gtg ctg aag ctg gtg Leu Ser Val Leu Arg Thr Phe Arg Leu Met Arg Val Leu Lys Leu Val 830 835 840 845	2907
cgc ttc ctg ccg gcg ctg cag cgg cag ctg gtg gtg ctc atg aag acc Arg Phe Leu Pro Ala Leu Gln Arg Gln Leu Val Val Leu Met Lys Thr 850 855 860	2955
atg gac aac gtg gcc acc ttc tgc atg ctg ctt atg ctc ttc atc ttc Met Asp Asn Val Ala Thr Phe Cys Met Leu Leu Met Leu Phe Ile Phe 865 870 875	3003
atc ttc agc atc ctg ggc atg cat ctc ttc ggc tgc aag ttt gcc tct Ile Phe Ser Ile Leu Gly Met His Leu Phe Gly Cys Lys Phe Ala Ser 880 885 890	3051
gag cgg gat ggg gac acc ctg cca gac cgg aag aat ttt gac tcc ttg Glu Arg Asp Gly Asp Thr Leu Pro Asp Arg Lys Asn Phe Asp Ser Leu 895 900 905	3099
ctc tgg gcc atc gtc act gtc ttt cag atc ctg acc cag gag gac tgg Leu Trp Ala Ile Val Thr Val Phe Gln Ile Leu Thr Gln Glu Asp Trp 910 915 920 925	3147
aac aaa gtc ctc tac aat ggt atg gcc tcc acg tcg tcc tgg gcg gcc Asn Lys Val Leu Tyr Asn Gly Met Ala Ser Thr Ser Ser Trp Ala Ala 930 935 940	3195
ctt tat ttc att gcc ctc atg acc ttc ggc aac tac gtg ctc ttc aat Leu Tyr Phe Ile Ala Leu Met Thr Phe Gly Asn Tyr Val Leu Phe Asn 945 950 955	3243
ttg ctg gtc gcc att ctg gtg gag ggc ttc cag gcg gag gga gat gcc Leu Leu Val Ala Ile Leu Val Glu Gly Phe Gln Ala Glu Gly Asp Ala 960 965 970	3291

17

aac aag tcc gaa tca gag ccc gat ttc ttc tca ccc agc ctg gat ggt	3339
Asn Lys Ser Glu Ser Glu Pro Asp Phe Phe Ser Pro Ser Leu Asp Gly	
975 980 985	
gat ggg gac agg aag aag tgc ttg gcc ttg gtg tcc ctg gga gag cac	3387
Asp Gly Asp Arg Lys Lys Cys Leu Ala Leu Val Ser Leu Gly Glu His	
990 995 1000 1005	
ccg gag ctg cgg aag agc ctg ctg ccg cct ctc atc atc cac acg gcc	3435
Pro Glu Leu Arg Lys Ser Leu Leu Pro Pro Leu Ile Ile His Thr Ala	
1010 1015 1020	
gcc aca ccc atg tgc ctg ccc aag agc acc agc acg ggc ctg ggc gag	3483
Ala Thr Pro Met Ser Leu Pro Lys Ser Thr Ser Thr Gly Leu Gly Glu	
1025 1030 1035	
gcg ctg ggc cct gcg tgc cgc cgc acc agc agc agc ggg tgc gca gag	3531
Ala Leu Gly Pro Ala Ser Arg Arg Thr Ser Ser Ser Gly Ser Ala Glu	
1040 1045 1050	
cct ggg gcg gcc cac gag atg aag tca ccg ccc agc gcc cgc agc tct	3579
Pro Gly Ala Ala His Glu Met Lys Ser Pro Pro Ser Ala Arg Ser Ser	
1055 1060 1065	
ccg cac agc ccc tgg agc gct gca agc agc tgg acc agc agg cgc tcc	3627
Pro His Ser Pro Trp Ser Ala Ala Ser Ser Trp Thr Ser Arg Arg Ser	
1070 1075 1080 1085	
agc cgg aac agc ctc ggc cgt gca ccc agc ctg aag cgg aga agc cca	3675
Ser Arg Asn Ser Leu Gly Arg Ala Pro Ser Leu Lys Arg Arg Ser Pro	
1090 1095 1100	
agt gga gag cgg cgg tcc ctg ttg tgc gga gaa ggc cag gag agc cag	3723
Ser Gly Glu Arg Arg Ser Leu Leu Ser Gly Glu Gly Gln Glu Ser Gln	
1105 1110 1115	
gat gaa gag gag agc tca gaa gag gag cgg gcc agc cct gcg ggc agt	3771
Asp Glu Glu Glu Ser Ser Glu Glu Glu Arg Ala Ser Pro Ala Gly Ser	
1120 1125 1130	
gac cat cgc cac agg ggg tcc ctg gag cgg gag gcc aag agt tcc ttt	3819
Asp His Arg His Arg Gly Ser Leu Glu Arg Glu Ala Lys Ser Ser Phe	
1135 1140 1145	
gac ctg cca gac aca ctg cag gtg cca ggg ctg cat cgc act gcc agt	3867
Asp Leu Pro Asp Thr Leu Gln Val Pro Gly Leu His Arg Thr Ala Ser	
1150 1155 1160 1165	
ggc cga ggg tct gct tct gag cac cag gac tgc aat ggc aag tgc gct	3915
Gly Arg Gly Ser Ala Ser Glu His Gln Asp Cys Asn Gly Lys Ser Ala	
1170 1175 1180	
tca ggg cgc ctg gcc cgg gcc ctg cgg cct gat gac ccc cca ctg gat	3963
Ser Gly Arg Leu Ala Arg Ala Leu Arg Pro Asp Asp Pro Pro Leu Asp	
1185 1190 1195	
ggg gat gac gcc gat gac gag ggc aac ctg	3993
Gly Asp Asp Ala Asp Asp Glu Gly Asn Leu	
1200 1205	

<210> 52
 <211> 1207
 <212> PRT
 <213> Artificial Sequence

<220>

<223> CACNA1G - a gene encoding a T-type calcium channel

<400> 52

```

Met Asp Glu Glu Asp Gly Ala Gly Ala Glu Glu Ser Gly Gln Pro
 1          5          10          15
Arg Ser Phe Met Arg Leu Asn Asp Leu Ser Gly Ala Gly Gly Arg Pro
 20          25          30
Gly Pro Gly Ser Ala Glu Lys Asp Pro Gly Ser Ala Asp Ser Glu Ala
 35          40          45
Glu Gly Leu Pro Tyr Pro Ala Leu Ala Pro Val Val Phe Phe Tyr Leu
 50          55          60
Ser Gln Asp Ser Arg Pro Arg Ser Trp Cys Leu Arg Thr Val Cys Asn
 65          70          75          80
Pro Trp Phe Glu Arg Ile Ser Met Leu Val Ile Leu Leu Asn Cys Val
 85          90          95
Thr Leu Gly Met Phe Arg Pro Cys Glu Asp Ile Ala Cys Asp Ser Gln
100          105          110
Arg Cys Arg Ile Leu Gln Ala Phe Asp Asp Phe Ile Phe Ala Phe Phe
115          120          125
Ala Val Glu Met Val Val Lys Met Val Ala Leu Gly Ile Phe Gly Lys
130          135          140
Lys Cys Tyr Leu Gly Asp Thr Trp Asn Arg Leu Asp Phe Phe Ile Val
145          150          155          160
Ile Ala Gly Met Leu Glu Tyr Ser Leu Asp Leu Gln Asn Val Ser Phe
165          170          175
Ser Ala Val Arg Thr Val Arg Val Leu Arg Pro Leu Arg Ala Ile Asn
180          185          190
Arg Val Pro Ser Met Arg Ile Leu Val Thr Leu Leu Leu Asp Thr Leu
195          200          205
Pro Met Leu Gly Asn Val Leu Leu Leu Cys Phe Phe Val Phe Phe Ile
210          215          220
Phe Gly Ile Val Gly Val Gln Leu Trp Ala Gly Leu Leu Arg Asn Arg
225          230          235          240
Cys Phe Leu Pro Glu Asn Phe Ser Leu Pro Leu Ser Val Asp Leu Glu
245          250          255
Arg Tyr Tyr Gln Thr Glu Asn Glu Asp Glu Ser Pro Phe Ile Cys Ser
260          265          270
Gln Pro Arg Glu Asn Gly Met Arg Ser Cys Arg Ser Val Pro Thr Leu
275          280          285
Arg Gly Asp Gly Gly Gly Gly Pro Pro Cys Gly Leu Asp Tyr Glu Ala
290          295          300
Tyr Asn Ser Ser Ser Asn Thr Thr Cys Val Asn Trp Asn Gln Tyr Tyr
305          310          315          320
Thr Asn Cys Ser Ala Gly Glu His Asn Pro Phe Lys Gly Ala Ile Asn
325          330          335
Phe Asp Asn Ile Gly Tyr Ala Trp Ile Ala Ile Phe Gln Val Ile Thr
340          345          350
Leu Glu Gly Trp Val Asp Ile Met Tyr Phe Val Met Asp Ala His Ser
355          360          365
Phe Tyr Asn Phe Ile Tyr Phe Ile Leu Leu Ile Ile Val Gly Ser Phe
370          375          380
Phe Met Ile Asn Leu Cys Leu Val Val Ile Ala Thr Gln Phe Ser Glu
385          390          395          400

```

Thr Lys Gln Arg Glu Ser Gln Leu Met Arg Glu Gln Arg Val Arg Phe
 405 410 415
 Leu Ser Asn Ala Ser Thr Leu Ala Ser Phe Ser Glu Pro Gly Ser Cys
 420 425 430
 Tyr Glu Glu Leu Leu Lys Tyr Leu Val Tyr Ile Leu Arg Lys Ala Ala
 435 440 445
 Arg Arg Leu Ala Gln Val Ser Arg Ala Ala Gly Val Arg Val Gly Leu
 450 455 460
 Leu Ser Ser Pro Ala Pro Leu Gly Gly Gln Glu Thr Gln Pro Ser Ser
 465 470 475 480
 Ser Cys Ser Arg Ser His Arg Arg Leu Ser Val His His Leu Val His
 485 490 495
 His His His His His His His Tyr His Leu Gly Asn Gly Thr Leu
 500 505 510
 Arg Ala Pro Arg Ala Ser Pro Glu Ile Gln Asp Arg Asp Ala Asn Gly
 515 520 525
 Ser Arg Arg Leu Met Leu Pro Pro Pro Ser Thr Pro Ala Leu Ser Gly
 530 535 540
 Ala Pro Pro Gly Gly Ala Glu Ser Val His Ser Phe Tyr His Ala Asp
 545 550 555 560
 Cys His Leu Glu Pro Val Arg Cys Gln Ala Pro Pro Pro Arg Ser Pro
 565 570 575
 Ser Glu Ala Ser Gly Arg Thr Val Gly Ser Gly Lys Val Tyr Pro Thr
 580 585 590
 Val His Thr Ser Pro Pro Pro Glu Thr Leu Lys Glu Lys Ala Leu Val
 595 600 605
 Glu Val Ala Ala Ser Ser Gly Pro Pro Thr Leu Thr Ser Leu Asn Ile
 610 615 620
 Pro Pro Gly Pro Tyr Ser Ser Met His Lys Leu Leu Glu Thr Gln Ser
 625 630 635 640
 Thr Gly Ala Cys Gln Ser Ser Cys Lys Ile Ser Ser Pro Cys Leu Lys
 645 650 655
 Ala Asp Ser Gly Ala Cys Gly Pro Asp Ser Cys Pro Tyr Cys Ala Arg
 660 665 670
 Ala Gly Ala Gly Glu Val Glu Leu Ala Asp Arg Glu Met Pro Asp Ser
 675 680 685
 Asp Ser Glu Ala Val Tyr Glu Phe Thr Gln Asp Ala Gln His Ser Asp
 690 695 700
 Leu Arg Asp Pro His Ser Arg Arg Gln Arg Ser Leu Gly Pro Asp Ala
 705 710 715 720
 Glu Pro Ser Ser Val Leu Ala Phe Trp Arg Leu Ile Cys Asp Thr Phe
 725 730 735
 Arg Lys Ile Val Asp Ser Lys Tyr Phe Gly Arg Gly Ile Met Ile Ala
 740 745 750
 Ile Leu Val Asn Thr Leu Ser Met Gly Ile Glu Tyr His Glu Gln Pro
 755 760 765
 Glu Glu Leu Thr Asn Ala Leu Glu Ile Ser Asn Ile Val Phe Thr Ser
 770 775 780
 Leu Phe Ala Leu Glu Met Leu Leu Lys Leu Leu Val Tyr Gly Pro Phe
 785 790 795 800
 Gly Tyr Ile Lys Asn Pro Tyr Asn Ile Phe Asp Gly Val Ile Val Val
 805 810 815
 Ile Ser Val Trp Glu Ile Val Gly Gln Gln Gly Gly Gly Leu Ser Val
 820 825 830
 Leu Arg Thr Phe Arg Leu Met Arg Val Leu Lys Leu Val Arg Phe Leu
 835 840 845
 Pro Ala Leu Gln Arg Gln Leu Val Val Leu Met Lys Thr Met Asp Asn
 850 855 860
 Val Ala Thr Phe Cys Met Leu Leu Met Leu Phe Ile Phe Ile Phe Ser
 865 870 875 880

20

Ile Leu Gly Met His Leu Phe Gly Cys Lys Phe Ala Ser Glu Arg Asp
 885 890 895
 Gly Asp Thr Leu Pro Asp Arg Lys Asn Phe Asp Ser Leu Leu Trp Ala
 900 905 910
 Ile Val Thr Val Phe Gln Ile Leu Thr Gln Glu Asp Trp Asn Lys Val
 915 920 925
 Leu Tyr Asn Gly Met Ala Ser Thr Ser Ser Trp Ala Ala Leu Tyr Phe
 930 935 940
 Ile Ala Leu Met Thr Phe Gly Asn Tyr Val Leu Phe Asn Leu Leu Val
 945 950 955 960
 Ala Ile Leu Val Glu Gly Phe Gln Ala Glu Gly Asp Ala Asn Lys Ser
 965 970 975
 Glu Ser Glu Pro Asp Phe Phe Ser Pro Ser Leu Asp Gly Asp Gly Asp
 980 985 990
 Arg Lys Lys Cys Leu Ala Leu Val Ser Leu Gly Glu His Pro Glu Leu
 995 1000 1005
 Arg Lys Ser Leu Leu Pro Pro Leu Ile Ile His Thr Ala Ala Thr Pro
 1010 1015 1020
 Met Ser Leu Pro Lys Ser Thr Ser Thr Gly Leu Gly Glu Ala Leu Gly
 1025 1030 1035 1040
 Pro Ala Ser Arg Arg Thr Ser Ser Ser Gly Ser Ala Glu Pro Gly Ala
 1045 1050 1055
 Ala His Glu Met Lys Ser Pro Pro Ser Ala Arg Ser Ser Pro His Ser
 1060 1065 1070
 Pro Trp Ser Ala Ala Ser Ser Trp Thr Ser Arg Arg Ser Ser Arg Asn
 1075 1080 1085
 Ser Leu Gly Arg Ala Pro Ser Leu Lys Arg Arg Ser Pro Ser Gly Glu
 1090 1095 1100
 Arg Arg Ser Leu Leu Ser Gly Glu Gly Gln Glu Ser Gln Asp Glu Glu
 1105 1110 1115 1120
 Glu Ser Ser Glu Glu Arg Ala Ser Pro Ala Gly Ser Asp His Arg
 1125 1130 1135
 His Arg Gly Ser Leu Glu Arg Glu Ala Lys Ser Ser Phe Asp Leu Pro
 1140 1145 1150
 Asp Thr Leu Gln Val Pro Gly Leu His Arg Thr Ala Ser Gly Arg Gly
 1155 1160 1165
 Ser Ala Ser Glu His Gln Asp Cys Asn Gly Lys Ser Ala Ser Gly Arg
 1170 1175 1180
 Leu Ala Arg Ala Leu Arg Pro Asp Asp Pro Pro Leu Asp Gly Asp Asp
 1185 1190 1195 1200
 Ala Asp Asp Glu Gly Asn Leu
 1205

<210> 53

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> primer for PCR (GAPDH)

<400> 53

cggagtcaac ggattggtcg tat

23

<210> 54

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> primer for PCR (GAPDH)

<400> 54
agccttctcc atggtggtga agac

24

<210> 55
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<400> 55
aaaaaaccca aactacaaaa ac

22

<210> 56
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 56
gttggtggrg ttgttggr

18

<210> 57
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 57
aactatcycc aacyccacaa

20

<210> 58
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 58
aagagatttt tttttttttt tttrgt

26

<210> 59
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> y = .C or T

<400> 59
aaaatccyaa aaaaaacycc ccc 23

<210> 60
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 60
ggaagtttta ggggrtagg ggaa 24

<210> 61
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 61
aacyatccct ccctctaacc tac 23

<210> 62
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<400> 62
aggtagtatg gtgaggtttg tttt 24

<210> 63
<211> 22
<212> DNA

<213> Artificial Sequence

<220>

<223> Target sequence for bisulfite-PCR primer

<221> misc_feature

<222> (0)...(0)

<223> r = G or A

<400> 63

atcaatacta aacraaatca aa

22

<210> 64

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Target sequence for bisulfite-PCR primer

<400> 64

aggaaaagaa aggtaaggg

19

<210> 65

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Target sequence for bisulfite-PCR primer

<221> misc_feature

<222> (0)...(0)

<223> r = G or A

<400> 65

caaaattaac rcaataaaaa aa

22

<210> 66

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Target sequence for bisulfite-PCR primer

<400> 66

tatttgaaga ggtggggaaa

20

<210> 67

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Target sequence for bisulfite-PCR primer

<400> 67

aaactcttac cccacctaac c

21

<210> 68
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 68
ggtttgtaat tggattaaay gtt

23

<210> 69
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<400> 69
ccactaactc aaaactaaaa aa

22

<210> 70
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<400> 70
gggaggtgta aaaggatgaa a

21

<210> 71
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<400> 71
ctaactactaa aataaaaata aa

22

<210> 72
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 72
gtaggatggt ataygaagag 20

<210> 73
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 73
aaacrctaac raacatacta c 21

<210> 74
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<400> 74
gggttttttt tagggtatatt 20

<210> 75
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 75
gaattaaatt tcaaaaaaac cr 22

<210> 76
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 76
tttaggagga tgyggagtt 19

<210> 77

<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 77
aaaaaaccta acraaacact ta 22

<210> 78
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<400> 78
gttattgtgt agtggagttt gg 22

<210> 79
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 79
actccratta acaaaccaac 20

<210> 80
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 80
aatatggttt yggttggtaa 20

<210> 81
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<400> 81
tccctaaatt ccacacatt 19

<210> 82
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<400> 82
gtaagttgta gttggttgtt tta 23

<210> 83
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 83
ctctctacta ccraattcct ct 22

<210> 84
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<400> 84
gttttggttt tggttgtg 18

<210> 85
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence for bisulfite-PCR primer

<400> 85
ccactaccaa acaaattcccc 20

<210> 86
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Target sequence for bisulfite-PCR primer

<400> 86

tttattgggg aatttcggg

19

<210> 87

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Target sequence

<221> misc_feature

<222> (0)...(0)

<223> r = G or A

<400> 87

aacaaaataa ctactacrcc rtc

23

<210> 88

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Target sequence

<400> 88

gtaaagtgag ggggtggtgat g

21

<210> 89

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Target sequence

<221> misc_feature

<222> (0)...(0)

<223> r = G or A

<400> 89

ctccaaaaaa ctataaatac ccraa

25

<210> 90

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Target sequence

<221> misc_feature

<222> (0)...(0)

<223> y = C or T

<400> 90

gagtgagtga aggyggtaga tt

22

<210> 91
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 91
aacctcacat taacrctcct aaa

23

<210> 92
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence

<400> 92
gttttttttaa gattgggttt ttttag

26

<210> 93
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence

<400> 93
caaaccctcaa acatccttta tcca

24

<210> 94
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence

<400> 94
ggatttaggg gtaaggggag gg

22

<210> 95
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 95
aaaaaccaca actaaatcc ratt

24

<210> 96
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence

<400> 96
agtgagggat ttagttgtgg tgtg

24

<210> 97
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 97
aactatcrcc aacrccacaa

20

<210> 98
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 98
aagagatttt tttttttttt ttttgt

26

<210> 99
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 99
aaaatccraa aaaaaacrcc ccc

23

<210> 100
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence

<221> misc_feature
<222> (0)...(0)
<223> y = C or T

<400> 100
ggaagtttta ggggygtagg ggaa

24

<210> 101
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence

<400> 101
aacaaaatac aactcccaaa caccc

25

<210> 102
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence

<400> 102
ttagggtttg attttttaat ttgggt

26

<210> 103
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence

<221> misc_feature
<222> (0)...(0)
<223> r = G or A

<400> 103
caaaaaatta cratcccccc tc

22

<210> 104
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Target sequence

<221> misc_feature
 <222> (0)...(0)
 <223> y = C or T

<400> 104
 ttggagggtat aataaggaga ttttygg

26

<210> 105
 <211> 576
 <212> DNA
 <213> Homo sapiens

<220>
 <221> gene
 <222> (0)...(0)
 <223> APOB CpG ISLAND

<400> 105
 cccgggaggc gcccttttga cctttttgcaa tcctggcgct cttgcagcct gggtcttcta 60
 taaatggggg gcgggcgcgc gccgcgcatt cccaccggga cctgcggggc tgagtgcctt 120
 tctcggttgc tgccgctgag gagcccgccc agccagccag ggccgcgagg ccgaggccag 180
 gccgcagccc aggagccgcc ccaccgcagc tggcgatgga cccgccgagg cccgcgctgc 240
 tggcgctgcc tgcgctgctg ctgctgctgc tggcgggcgc cagggccggg gaggcgcgcg 300
 ccgctctgcg ggcagcagag ggagcgggag ggagccggcg gaccgagggt ggccggggca 360
 gcctgggcct aggccagagg gagggcagcc acaggggtcca gggcgagtgg ggggattgga 420
 ccagctggcg gcccctgcag gctcaggatg gggggcgcgg gatggagggg ctgaggaggg 480
 ggtctccgga gcctgcctcc ctctgaaag gtgaaacctg tgccgggtgt cccctgtcg 540
 ggcccctagc acccgctggg aagacgtggg aagctc 576

<210> 106
 <211> 2093
 <212> DNA
 <213> Homo sapiens

<220>
 <221> gene
 <222> (0)...(0)
 <223> CACNA1G CpG ISLAND

<400> 106
 cctcgggccc tacgccagga ccccgcgccg aatactctga ttcttcgggc tccctccaag 60
 ggagtcccaa agaccccaat ggccaatagg aaagtgggtt cggctctggc agcagtctga 120
 ttggctccag ccttcgggag cggaccagag ggcaagggga ggggagaggg gcggtcctgg 180
 gttttggggg gggaatcgga ttccagctgt ggttctctcc ctgcgctccc gcccgactg 240
 ccacggcgga cggccaatgg gcgcgcggct cggggccggc ggctccggc gattggctgc 300
 ggggctgtct gggggcgggg ccgaggcttg aagttagagt gagggatcca gctgtggtgt 360
 gcgcggggct cctcgccgcc gctttcgctc gctcgctccg cgtctcgccc ggagaggag 420
 gctgtggcgc cggcgacagc tacggcagcg gcagccaccg cggcggtgc ggcgccgga 480
 tctccgctc cactcccgcc cgggactgcc cccactgtc tccccgccc tcccgagacag 540
 tgagcccgcg gcggggcggg ggaaggagcc gccccaccc cctccaagcc caccctaaa 600
 gagatccctc ctccctccc ccgcccctg gcgcggagcc gggacgatgc tgaccctta 660
 gatccggtc cagctgcgcc gcgggaagag ggggcgcccc tccccgacc cccgccctcc 720
 gccgtgccc ccttttctg tgcctctctc ggggcggctt cgcgaaggat agcgccgaat 780
 ccggcaaccg gagcctgggc gcgaagcgaa gaagccggaa caaagttagg gggagccggc 840
 cggctggccc ggggaagcccc aggggcgcag ggaagcggg actcgcgcg ggcggggttt 900
 cctgcgccc cggcgccccg cgggcagcat gccctgcgg gcagggggag ctgggctgaa 960
 ctggccctcc cgggggctca gcttgccccc tagagccac cagatgtgcc ccgcggggg 1020
 ccccggggtt gcgtgaggac acctcctctg aggggcgcgg cttgcccctc tccggatcgc 1080
 ccggggcccc ggctggccag aggatggacg agggaggagga tggagcgggc gccgaggagt 1140
 cgggacagcc ccggagcttc atgcggctca acgacctgtc gggggccggg ggcgccggg 1200

gccgggggtca	gcagaaaagg	acccgggcag	cgcggtactcc	gaggcggagg	ggctgccgta	1260
cccggcgctg	gccccggtgg	ttttcttcta	cttgagccag	gacagccgcc	cgcgagctg	1320
gtgtctccgc	acggtctgta	acccatatcc	ttcggggcac	gacggccagg	cgcggggtca	1380
gaagggggac	gggcccgcacc	gccgggggtc	gggggggaag	aagaccacc	gccaggtgag	1440
tcgaagtgag	cccggagggt	aggcggtatg	ggggggggct	gccagggagg	ggagggggca	1500
ccagagtggg	agcgagagcg	cgagcaggtc	tcgtcggtaa	cccgggctta	ccccacctgc	1560
gtacacacac	ctcagtcctc	ctgggttggg	ggggtgggga	tccaggccag	gagaagagag	1620
ctgtgccccg	ctggctcgca	gctggacgcc	ctccagatgt	ggtcagggga	gggtcgtcat	1680
cctccagatg	tgggaagctt	cgggagcctg	ggagctgtac	tctgcccgcg	ccggttagcg	1740
agctgggttt	ggtttccgag	tttgggtggg	ggtgggtggg	ggcgggtggg	aggaagctgc	1800
ggggacggag	gaggggggac	cgcaatctcc	tgggtttccc	tccttcccc	gccccaaagt	1860
ttgcggcgga	ttctagatgt	tggggggcgg	ggaccaggtc	ctggcccacc	tcacccccca	1920
cctcgcggtg	tggaggcaca	acaaggagat	tccggcgcg	gctgatgtca	ggggcgcgaga	1980
atgagaacaa	gatgtgggtg	aggggagctg	tctgcccccg	gagctgggag	tggagccctt	2040
ttccgctaga	gcccagtgcc	gcgggtgcct	cctacccgat	ctccattcga	tgc	2093

<210> 107

<211> 327

<212> DNA

<213> Homo sapiens

<220>

<221> gene

<222> (0)...(0)

<223> CDX2 CpG Island

<400> 107

ctcgtaaatc	acggaagccg	ccggcctggg	gctccgcacg	ccagcctgtg	cggtctcttc	60
ccgcctctgc	agcctagtgg	gaaggagggt	ggaggaaaga	aggaagaaag	ggagggaggg	120
aggaggcagg	ccagagggag	ggaccgcctc	ggaggcagaa	gagccgcgag	gagccagcgg	180
agcaccgcgg	gctggggcgc	agccaccgcg	cgctcctcga	gtccccctcg	ccctttccct	240
tcgtgcccc	cggcagcctc	cagcgctcgt	ccccaggcag	catggtgagg	tctgctcccc	300
gtccctcgcc	accatgtacg	tgagcta				327

<210> 108

<211> 1663

<212> DNA

<213> Homo sapiens

<220>

<221> gene

<222> (0)...(0)

<223> EGFR CpG Island

<400> 108

gtccgcgggg	accgggtcca	gaggggcagt	gctgggaacg	cccctctcgg	aaattaactc	60
ctcagggcac	cgctccccct	ccatgcgccg	ccccactccc	gccggagact	aggtcccgcg	120
ggggccaccg	tgteccaccg	ctcgcgccg	ctggccttgg	gtccccgctg	ctgggtctctc	180
tcctctctcc	tcgcattctc	ctcctcctct	gtcctctccg	atccctcctc	cgccgcctgg	240
tcctctctcc	tcctcgccctg	cctcccgccg	ctcgggcccg	gcgagctaga	cgctccggga	300
gcccccgccg	cagcgcgccg	gcagcagcct	cctccccccg	cacggtgtga	gcgcccgcgg	360
cgcgaggcgg	gccggagtcc	cgagctagcc	ccgcggcccg	cgccgcccag	accggacgac	420
aggccacctc	gtcgcgctcg	cccagatccc	cgctctcgcc	ccaacgccac	aaccaccgcg	480
cacggccccc	tgactccgtc	cagtattgat	cgggagagcc	ggagcgagct	cttcggggag	540
cagcgatgcc	accctccggg	acggccgggg	cagcgctcct	ggcgctgctg	gctgcgctct	600
gccccgcgag	tcgggtctcg	gaggaaaaga	aaggtaaggg	cgtgtctcgc	ggctccccgc	660
cgcccccgga	tcgcgccccg	gacccccgag	cccgcccaac	cgcaccgcgc	accggtctcg	720
cccgcgcccc	cgcccgctct	ttctgttttc	cttgagatca	cgtgcgcccg	cgaccgggac	780
cgcgggagga	acgggacggt	tcgttctctg	gccgggagag	tctggggcgg	gcggaggagg	840
agacgcgtgg	gacaccgggc	tgcaggccag	gcggggaaacg	gccgccggga	cctccggcgc	900

```

cccgaaaccgc tcccaacttt cttccctcac tttccccgcc cagctgcgca ggatcggcgt      960
cagtggggcga aagccgggtg ctggtgggcg cctggggccg ggggtcccga cgggctcccc      1020
gcgctgtctt cccagggcgc gacgggggtc tggcgcgcac ccgagggccg ctgcccaccc      1080
gccgagactg cctgtttagg gaagctgagg aaggaaccca aaaatacagc ctccgcctcg      1140
accccgcggg acagggcggt ttctgagagg acctccccgc ctccgcgctc cgcgcaggtc      1200
tcaaaactgaa gccggcgccc gccagcctgg ccccgcccc tctccaggtc cccgcgatcc      1260
tcgttcccca gtgtggagtc gcagcctcga cctgggagct gggagaactc gtctaccacc      1320
acctgcggct cccggggagg ggtggtgctg gcggcggtta gtttcctcgt tggcaaaagg      1380
caggtggggg ccgaccgcgc ccttgggcgc agaccgcggc cgctcgctc gcccggtgcg      1440
ccctcgtctt gcctatccaa gagtgcctcc cactcccggg accccagctc cctccgcgcc      1500
cgcgccgaaa gccccaggct ctccctcgat ggccgcctcg cggagacgtc cgggtctgct      1560
ccacctgcag cccttcgggtc gcgcctgggc ttcgcggtgg agcgggacgc ggctgtccgg      1620
ccaactgcagg gggggatcgc gggactcttg agcggaagcc ccg                        1663

```

<210> 109

<211> 1787

<212> DNA

<213> Homo sapiens

<220>

<221> gene

<222> (0)...(0)

<223> FBN1 CpG Island

<221> misc_feature

<222> (1)...(1787)

<223> n = A,T,C or G

<400> 109

```

agagccgcgt ctggagtggt ctctcgacac ccagggcaag tgggggcggc agagccctct      60
cctcggtcgg cacagcagcc tctgcgcgg tcccggcctg cgacgcgccc agtcttagcc      120
tcccggcctc cggcgtctgc tgagtgtccg gcgggagagg cgaggaggag gcgctaccgg      180
gagggcgggg cagcgggggac tgggtttctc tcggggccagg gcctccgggg caaccgtctc      240
cagcgcgcat tcttggtgca ggtggaacag cttctgtctc cggtagggct tcacctatcg      300
cgggagaggt taatctcgga tctaaacctc gcagcgcgag agcgggctaa aaccgctact      360
ccacctcttc ccatttctcc cctccccacc tcaagacaaa aagtcccagg ccgggcaggga      420
cctgatcacc tctgcctcct cccactgcgc taatcctgcg agcgagaggc cccgcaccga      480
ggcggagggt gcaaagggga gtggaaaggg aggatggatg gggccggggg gtggggtggt      540
gatgagggcg acgaaggagg ggggtgtcatt ttctttttct ttcttttttt aaaaaagta      600
tttctctcgc gagaaccgc tgccgggacg atacttgaag aggtggggaa aggagggggc      660
tgccggagcc gcggcagaga ctgtgggtgc cacaagcgga caggagccac agctgggaca      720
gctgcgagcg gagccgagca gtggctgtag cggccacgac tgggagcagc cgcgcgcgcc      780
tcctcgggag tcggagccgc cgcttctcca gtgggtgcag ccgggggtccg acgggggtcg      840
ggcggccacc ggggctggag ctgcggccac ggaggctttt gcgtttgcgc cgnnngaggg      900
cagggacagg gactgggggt aggggctgtc ccggaacgtc caacgtggnc gctggacctt      960
cccctgcctg acagcttctc gnccgggggt tcttgggtgc ggnccggcgt cagatgttcg      1020
ggggggcggt catcgcccgg agtcggcggg gacggcgcg ctgcttccag ctggcggaga      1080
gggcaggctg aggagtgggg cgttcagagc gcgcacgcg cgcaattcgt gccgctaaaa      1140
aaaataaacc cagagagctc gcccggggct taggaccgct ggggatatgg gtactttgcg      1200
ccgcgctctt ctggcggggc ccgggaggcg agggattggc cggggctgct gcgccggggc      1260
ctgggctttc cagccagctg tggaccaaac ggtcttccct taccctaaatt aactgcgcca      1320
caggcgcccg acnggttggg ctttgggaat ggggaccgcg agcttcagca tcccgatgcc      1380
ctgaaagtct ccccgccctc gggatttgtc tctgtgttgc agctggcagg ggcgcgctga      1440
agtgggagca gcgctggag aaggcgggag gagcccgccc cgggggacgg gcggcgggat      1500
agcgggaccc cgcggcgcg gtgcgcttca gggcgagcg gcggccgag accgagcccc      1560
gggcgcggca agaggcgcg ggagccggtg gcggctcgcc atcatgcgtc gagggcgctc      1620
gctggagatc gccctgggat ttaccgtgct tttagcgtcc tacacgagcc atggggcgga      1680
cgccaatttg gaggctggga acgtgaagga aaccagagcc agtcggggcca agagaagagg      1740
cgggtggagga cagcagcgcg ttaaagggta aaggaaccgg ttccctc                        1787

```

<210> 110
 <211> 810
 <212> DNA
 <213> Homo sapiens

<220>
 <221> gene
 <222> (0)...(0)
 <223> GPR37 CpG Island

<400> 110
 tccccgcccc caccgcgccc tagccccgggc tcgggggacct gtcagggtgg tttcgacagc 60
 tgggggaatta acctgtcccc cccatcccta gcctcgagcc gcgcaggctc cgcgcctccg 120
 cccttggttc ctcccagctc ctcccgagtgg aagccgctac aaatggcttg aatgaaacgt 180
 gtgtgggttt agtgagtggg gaaccaccag gggatcccggt cccccacaa accagtatct 240
 ctccgaggag gaggcgaagg agtgggagga ggcaacgagc cgagagtcca gcttcgcggg 300
 cgcgcgcagc ggctggagcg cgggggagcg gccgggcccac ctccccttcc cggccgcgca 360
 ctgcctggcc cgcggcggtt ccaggcacca ccttcccgt cgggctgag cccgctgtgg 420
 cagtgactag ctcccgcggc tagcggcact gtccaccgac gagcggcgcc ctcttctccc 480
 ccttctcccc acgatttctt tctctgcggc ggcacgcct ccagcagcct gcttcgcccc 540
 gtctgtaact ttgagctgga ggagaagcaa ctttggcagt ggccgcgggg ttggaatccc 600
 gcttctcttc ggcagcagta ggctcgcaag tcgctggggg taggtggggc aagagtttcg 660
 ccggcgcacg agcgtgtctt cggactgttt gcaacgtgtt tccagcgagc tgggagcggg 720
 gttgtgactg cgagtcgtct gggggagggg gacttggttt tcttttcttc tagagacctc 780
 ggcttgcaac tggatcaaac gctgtcgaaa 810

<210> 111
 <211> 550
 <212> DNA
 <213> Homo sapiens

<220>
 <221> gene
 <222> (0)...(0)
 <223> HSPA6 CpG Island

<400> 111
 tgtatttcgca tggtaacata tcttcgggtct tcctgccgct gggctctcag cggccctcca 60
 aggcagcccc caggccccgtg ctgcctcag ggatccctcca cagccccggg gagaccttgc 120
 ctctaaagtt gctgcttttg cagctctgcc acaaccgcgc gtcctcagag ccagccggga 180
 ggagctagaa ccttccccgc gtttctttca gcagccctga gtcagaggcg ggctggcctt 240
 gcaagtagcc gcccagcctt cttcgggtctc acggaccgat ccgcccgaac cttctccggg 300
 ggtcagcgcc gcgctgcgcc gcccggtga ctcagcccgg gcgggggggc gggaggtct 360
 cgactggggc ggaaggtgcg ggaaggttcg cggcgggcgg gtcggggagg tgcaaaagga 420
 tgaaaagccc gtggacggag ctgagcagat ccggccgggc tggcggcaga gaaaccgcag 480
 ggagagcctc actgctgagc gccctcgcac gcggggcgga gcagcctccg tggcctccag 540
 catccgacaa 550

<210> 112
 <211> 278
 <212> DNA
 <213> Homo sapiens

<220>
 <221> gene
 <222> (0)...(0)
 <223> IQGAP2 CpG Island

<400> 112
 agagttcact tttacttcag tgtcagcgcg cggcgggcgt ggctgggtct ggcgagagag 60

36

caccgagggga	gtgggtcgca	gatcttcggg	cggctagggg	aaatcggcga	gaggcgggat	120
ccgagcgcg	cggcgggcg	cagagccgc	gagcctggc	agcgagggt	gccgcgggg	180
gcgcgcccc	ggcgggccc	cggagacgc	caggatgcca	cacgaagag	tgccgtcgct	240
gcagagaccc	cgctatggc	ctattgtgga	cgatgaaa			278

<210> 113
 <211> 1461
 <212> DNA
 <213> Homo sapiens

<220>
 <221> gene
 <222> (0)...(0)
 <223> KL CpG Island

<400> 113						
ctcgaaagag	gggcgcgggt	gggcgcgtct	ccccgcgagc	atctcaccta	agggggaatc	60
cctttcagcg	cacggcgaa	ttccccctcg	gctgtccac	ctggcagtc	ctctaggatt	120
tgggccagtc	cctaattggc	tccagcaatg	tccagccgga	gcttctttgg	gcctccgagt	180
gggagaaaag	tgagagcagg	tgcttcccc	gcggcgcgct	ccgctagggc	ccggcaggat	240
ccgcgcccc	agtcgggaaa	agttgggtcg	cgccttttct	ccccgacgaa	gccgtccag	300
ggctgctctc	agaggacgcg	cggcaggcaa	agagaatgaa	cctgagcgtc	cacgaaacgt	360
cctgcacggc	tcccgggagc	tgaggagaa	aggtgccttt	ctccgacgtc	cgcgggcgac	420
gcctgcgcga	ccttgcgccg	tgccgcgccc	ctcccgggca	cccctcgccc	tcggcgcccc	480
tgccccacc	cccagtgcca	gggcggaggc	agtcgccggt	cgcaggtaat	tattgccagc	540
ggagcccgcc	ggggagcggg	ggtgggcgcg	cgggcgggtg	gcgggcgggc	gcggcggggc	600
gcgggcataa	aggggcgcgg	cgcggggccc	cggagcctgg	ctcccgcgca	gcatgcccgc	660
cagcgccccg	ccgcgcgcgc	cgcggccgcc	ggcgccgtcg	ctgtcgctgc	tgctggtgct	720
gctgggcctg	ggcgccgcgc	gcctgcgtgc	ggagccgggc	gacggcgcg	agacctgggc	780
ccgtttctcg	cggcctcctg	cccccgaggc	cgcgggcctc	ttccagggca	ccttccccga	840
cggcttctc	tgggccgtgg	gcagcgccgc	ctaccagacc	gagggcggct	ggcagcagca	900
cggcaagggt	gcgtccatct	gggacacgtt	caccaccac	cccctggcac	ccccgggaga	960
ctcccggaa	gccagtctgc	cgttgggcgc	cccgctcgcc	ctgcagccc	ccaccgggga	1020
cgtagccagc	gacagctaca	acaacgtctt	ccgcgacacg	gaggcgctgc	gcgagctcgg	1080
ggtcactcac	taccgcttct	ccatctcgtg	ggcgcgagtg	ctcccgaatg	gcagcgcggg	1140
cgtccccaac	cgcgaggggc	tgcgctacta	ccggcgccctg	ctggagcggc	tgcgggagct	1200
gggcgtgcag	cccgtggtca	ccctgtacca	ctgggacctg	ccccagcgcc	tgaggagcgc	1260
ctacggcggc	tgggccaacc	gcgccctggc	cgaccacttc	agggattacg	cggagctctg	1320
cttcggccac	ttcggcggtc	aggtcaagta	ctggatcacc	atcgacaacc	cctacgtggt	1380
ggcctggcac	ggctacgcca	cgggcgcct	ggccccggc	atccggggca	gcccgcggct	1440
cgggtacctg	gtggcgca	a				1461

<210> 114
 <211> 249
 <212> DNA
 <213> Homo sapiens

<220>
 <221> gene
 <222> (0)...(0)
 <223> PAR2 CpG Island

<400> 114						
cccggggcgt	ggcctccgc	aggtgagtac	gctgctcctt	cggtttccct	gaaacctaac	60
ccgcccggg	gaggcgcgca	gcagaggctc	cgaatcgggg	caggtgagag	gctgactttc	120
tctcggtgcg	tccagtggag	ctctgagttt	cgaatcgggc	gcggcggtat	ccccgcgcg	180
ccggcgctcg	ggcttccagg	aggatgcgga	gccccagcgc	ggcggtggctg	ctggggggccg	240
ccatcctgc						249

<210> 115

<211> 709
<212> DNA
<213> Homo sapiens

<220>
<221> gene
<222> (0)...(0)
<223> PITX2 CpG Island

<400> 115
agtcctgtgct cctgtctctc gggtgggtcc taagtgtccc gccaggtccc ctctcctttc 60
gctctcccgg ctccgggtcc cgactcttcg gcccgctggc atctgtcttc ctcccctgcc 120
tcgtttctcg tcgccccctgc tcgtcccccc cggcgctcgc ccgggcgctg tgctcgctcc 180
tggatcgcca gccgcgcagc cgggctcggc cggccgcccg cgcgccactg tgcagtggag 240
tttgggtgaa tctctgtctga cgtcacgtca ctccccacac ggagtaggag cagaggggaag 300
agagagggat gagagggagg gagaggagag agagtgcgag accgagcgag aaagctggag 360
aggagcagaa agaaactgcc agtggcggct agatttcgga ggccccagtg caccctgtga 420
ctccttcgga acttggcacc ctccaggagcc ctgcagtcct ctccaggccc gctttcgggc 480
gcttgccgtg cagccggagg ctccgctcgc tggaaatcgc cccgggaagc agtgggacgc 540
ggagacagca gctctctccc ggtagccgat aacggggaaa tggagaccaa ctgccgcaaa 600
ctgggtgcgg cgtgtctgca attagagaaa gataaaagcc agcaggggaa gaatgaggac 660
gtgggcgcgg aggaccgcgc taagaagaag cggcaaaagg gccagcgga 709

<210> 116
<211> 1496
<212> DNA
<213> Homo sapiens

<220>
<221> gene
<222> (0)...(0)
<223> PTCA CpG Island

<221> misc_feature
<222> (1)...(1496)
<223> n = A,T,C or G

<400> 116
gcggccgcag cggcagcagc gcccgccgtg tgagcagcag cagcggtctg tctgtcaacc 60
ggagcccag cccgagcagc ctgcggccag cagcgtcttc gcaagccgag cgcccaggcg 120
cgccaggagc ccgcagcagc ggcagcagcg cgcggggccg cccgggaagc ctccgtcccc 180
gcggcgccgg cggcgccgag ggcaacatgg cctcggctgg taacgccgcc gagccccagg 240
accgcggcgg cggcgccagc ggctgtatcg gtgccccggg acggccggct ggaggcggga 300
ggcgagacg gacggggggg ctgcgccgtg ctgcgcgcc ggaccgggac tatctgcacc 360
ggcccagcta ctgcgacgcc gccttcgntn nggagnagat tnccanggn nggcatttca 420
gactntntcn ttcccacttt ntcttccent acctntaact cntnggggat cgccccgcgc 480
acacacaaac acacacactn tcttctctn tntctcacac acaacacaca cactcactca 540
cacntctnca ggaaaagcag cagacaaatg gggattgaaa aattcaaacc ctccctctgg 600
tnntgggagg aaagggctgt ctgaggtccg cagggggtgg aggtgtgtgt gtgtgcgtgt 660
gtgtgtgtgn anacacacgc cctccctggg gtgccttttc cggagcactg gaaagccgtc 720
cacggcggac cacctcaagg gcggccgcgc ggtcgtagcg gtagtagcgt tcgtcgtgtg 780
agtagtagta gcggttggtt tgtaaatcgg agttcgagtt cgagtagttt gcggttagta 840
gcgttttcgt aagtcgagcg ttaggcgcg ttaggagttc gtagtagcgg tagtagcgcg 900
tcgggtcggt cgggaagttt tcgttttcgc ggcgccggcg gcggcgccgg taatatggtt 960
tcggttggtt acgtcgtcga gttttaggat cgcggcggcg gcggtagcgg ttgtatcggt 1020
gtttcgggac ggtcgggttg aggcgggagg cgtagacgga cggggggggt gcgtcgtgtt 1080
gtcgcgtcgg atcgggatta tttgtatcgg ttagttatt gcgacgtcgt tttcgtntng 1140
gagnagattt nttanggnng gtattttaga ttntntntnt tttattttnt tttttntat 1200
ttntaattn tnggggatcg ttttcgttat atataaatat atataatntt tttttntntn 1260
ttttatatat aatatatata tttatttata tntttntagg aaaagtagta gataaatggg 1320

gattgaaaaa	tttaaatttt	ttttttggtt	ntgggaggaa	agggttgttt	gaggttcgta	1380
gggggtggag	gtgtgtgtgt	gtgcgtgtgt	gtgtgtgnan	atatacgttt	tttttggtgt	1440
gtttttttcg	gagtattgga	aagtcgttta	cggcggatta	ttttaagggc	ggtcgt	1496

<210> 117

<211> 701

<212> DNA

<213> Homo sapiens

<220>

<221> gene

<222> (0)...(0)

<223> PTCHB CpG Island

<221> misc_feature

<222> (1)...(701)

<223> n = A,T,C, or G

<400> 117

gcggccgcgg	cactgtcctg	ccccgtgccc	cctgccctga	actttcttct	cctgcgcccc	60
tgccccctatt	tgacgcctaa	actcctgtac	ggctgccaca	tttcttaaca	tcttgagggg	120
ggaggcggag	tggagagagg	cggagagagg	aaggggggag	ggagccgaaa	taaagggtgt	180
ttcctttttt	ggcagccagt	tttggttttg	ttgagcatga	aatctctgct	cccttaaaaa	240
attattctcg	gaaaaagata	tccccccgt	tttccaggtt	ttgagccgcc	tctccttagg	300
gcctggtcgg	gggaggaaaa	gttgtaaaca	aattgccacc	ttaaattcgc	ggtgcgantic	360
tgcgagactg	ccgggttcat	tgtgttttac	aggctcgtcg	aaatgtgtgg	aatccagggg	420
aggcgagcac	ccagacgggg	gcccgcgggg	gccgcggcca	gcgcggggga	aatgccgcgc	480
cggggagcag	catgcgccgg	cctgagccct	tccctttgca	ctcggtctgt	ttttacgttt	540
aaccagaaag	gaaggagag	gagggaaaga	tccatgtggc	tgccctcttc	cgatcacaaa	600
tattgtcgta	agttgcagct	ggctgcccc	nttctaatt	cagctcacac	agcntntccc	660
cacgctatgg	aaatgcgtcg	ggagtgaact	ccggcgccgg	c		701

<210> 118

<211> 273

<212> DNA

<213> Homo sapiens

<220>

<221> gene

<222> (0)...(0)

<223> SDC1 CpG Island

<400> 118

ggagaggtgc	gggccgaatc	cgagccgagc	gagaggaatc	cggcagtaga	gagcggactc	60
cagccggcgg	accctgcagc	cctcgcctgg	gacagcggcg	cgctgggcag	gcgcccaga	120
gagcatcgag	cagcgaacc	cgcaagccg	gcccgcagcc	gcgacccgcg	cagcctgccg	180
ctctcccggc	gccggtccgg	gcagcatgag	gcgcgcggcg	ctctggctct	ggctgtgcgc	240
gctggcgctg	agcctgcagc	tgccctgcc	gca			273

<210> 119

<211> 751

<212> DNA

<213> Homo sapiens

<220>

<221> gene

<222> (0)...(0)

<223> SDC4 CpG Island

<400> 119

39

```

agtaggagcc ggcgggctcg ggcagggcgg gtcccttggg gtttccaact ccgcgggcgg      60
gcgcagtgcc ccgcaggcct cgcttccact ggggaattcc gggcggggtg cggcgggcgg      120
ggcgggggcg ggccggggcg gggccggtag gccgcctata agatgggtgg cgcgcccgcc      180
cggggccact cgccgcagcc tgcgcgcctt ctccagtcgg cggtgccatg gcccccgccc      240
gtctgttcgc gctgctgctg ttcttcgtag gcggagtcgc cgagtcggtg ggtgcttgga      300
ggttcccggg ctgggggcga agcggggggc caggccgggt cctcctttgt tcgtcggagc      360
gtgggatggg ggggggcaga tcgggggtac gctaccccca accggacacc gagggccggg      420
aaactttggt ggaaactttg ctccggggtc acgggccagc ctccgggatg gcttcacgcg      480
ccgtgcgccc ctgcctgtt gctcttccc cctccccggg cctcagcccc gcccgggcct      540
acgggctcgt tagtgactaa gccggtgtca actcttcaac tcccacacc tcgtcccttc      600
cctggtgacc ctggggcagg ctggagcgc tgaatccct cctcgtcttc ggggcgcccc      660
gagcagacag ctttaggatc cgagatggcc ctgggggtcg gggggctgcg tgtactcgga      720
agggggaggg ttttaggggt gtgcgaggcc c                                751

```

<210> 120

<211> 673

<212> DNA

<213> Artificial Sequence

<220>

<223> DNA fragment termed MINT31

<221> misc_feature

<222> (1)...(673)

<223> n = A,T,C or G

<400> 120

```

cccggggcct ctatcctggc gggaagggca ggccgaccgg gcagactgcg gcctctcggg      60
aggaagaag gtgtcagacg cgcggagcaa ccataaatag ccccccttcc ccagaagacg      120
gcacgggggt caagactcag gcgcgcgata ctcagaatga gagcagagac tcccgccagg      180
aaaaaagggc acttagggga tctgctcatt aacatgaaat gcaaatgagc ccgcccggcc      240
tcattttacac aactctgtgc atggattcgg cgaaagggca accagggaga cgacggcgca      300
gcagccactc tgccacttcc cccatccctt cccccccatc ggccggggcg ggaactgaga      360
cgaccccaac cctctgcggc ggcgggaggt gcgcgggggc tgcgtgggtg gtgcagcctt      420
aggggagtg acaacgcccc ggggtgatgg cctcagcaaa gtgaggggtg gtgatggagg      480
tcattccgacc catcccgccg cctctccgca gtggcgcaag cggcccaaaa tctccggaga      540
nggaactgag tgaccacta gggtccgccc tgtctacctc tcgcagatgt tggggaagtg      600
cttcccggcg tctaatectc gctgttcccc cctccaccgg cgcccagcac acccgcggcg      660
ctccgctccc ggg

```

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/25479

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07H 21/04; C12Q 1/68

US CL : 536/23.1; 435/6

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.1; 435/6

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,552,277 A (NELSON et al) 03 September 1996, see entire document.	10-11, 13, 19, 22-24
X	US 5,756,668 A (BAYLIN et al) 26 May 1998, see entire document.	10-11, 13, 19, 22-24
X	BAYLIN et al. Alterations in DNA Methylation: A Fundamental Aspect of Neoplasia. Advances in Cancer Research. 1998, Vol 72, pages 141-196, see entire document.	10-11, 13, 22-24
X	Database Medline US Library of Medicine, AN AF124351, TOYOTA et al., June 1999.	1-9, 31-32
—	'Inactivation of CACNA1G, a T-type calcium channel gene, by aberrant methylation of its 5' CpG island in human tumors'.	25-27
Y	Database Genbank on STN, AN AF027984, PEREZ-REYES et al. October 1998	1-2, 5, 31
X	'Molecular characterization of a neuronal low voltage-activated, T-type calcium channel'.	25-27
—	Nature. 391, 8896-896 (1998).	
Y	PEREZ-REYES et al. Molecular Characterization of Two Members of the T-Type Calcium Channel Family. Annals of the New York Academy of Sciences. 30 April 1999, Vol 868, pages 131-143.	1-9, 31-32
A	PEREZ-REYES et al. Molecular Characterization of a neuronal low-voltage-activated T-type calcium channel. Nature. 26 February 1998, Vol 391, pg 896-900.	1-9, 31-32

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "B" earlier application or patent published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

23 October 2000 (23.10.2000)

Date of mailing of the international search report

28 NOV. 2000

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks

Box PCT

Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Jeanine Enewold Goldberg

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/25479

Continuation of B. FIELDS SEARCHED Item 3: Medline, Biosis, Caplus, Embase, Scisearch
hypermethylation, CACNA1G, cancer, list of genes